

On some remarkable operads constructed from Baxter operators ¹

Philippe LEROUX

Institut de Recherche Mathématique, Université de Rennes I and CNRS UMR 6625
Campus de Beaulieu, 35042 Rennes Cedex, France, pleroux@univ-rennes1.fr

20/10/03

Abstract: Language theory, symbolic dynamics, modelisation of viral insertion into the genetic code of a host cell motivate the introduction of new types of bialgebras, whose coalgebra parts are not necessarily coassociative. Similarly, certain algebraic descriptions of combinatorial objects such as weighted directed graphs through a coalgebraic formulation often require at least two coalgebras whose coproducts verify entanglement conditions. All these structures generate special types of associative algebras (or binary quadratic and non-symmetric operads) obtained from commuting Baxter operators. One of the aim of this article is to study what type of associative algebras appear when such or such coalgebraic structures are used to describe for instance combinatorial objects. Examples from symbolic dynamics, weighted directed graphs and matrix theory are given.

Contents

1	Introduction and notation	2
2	Algebraic structures related to one co-operation	6
2.1	$\epsilon'[R]$ -Bialgebras and L -anti-dipterous algebras	6
2.2	On $\epsilon[R]$ -bialgebras	8
2.3	$\epsilon'[L]$ -Bialgebras and L -anti-dipterous algebras	14
2.4	On $\epsilon[L]$ -bialgebras	16
2.5	ϵ -Bialgebras	18
2.6	Construction of ϵ -bialgebras from weighted directed graphs	19
2.7	Other constructions	21

¹2000 *Mathematics Subject Classification*: 17A30, 18D50. *Key words and phrases*: dendriform dialgebras, quadri-algebras, octo-algebras, dipterous algebras, coassociative codialgebras, coassociative cotrialgebras, Leibniz algebras, Poisson dialgebras, Baxter operators.

3	Algebraic structures related to two entangled coproducts.	22
3.1	L -Coalgebras $(A, \mu, \Delta_1, \Delta_2)$, with (A, μ, Δ_2) a $\epsilon'[R]$ -bialgebra.	22
3.2	Dendriform dialgebras from L -coalgebras	24
3.3	Entanglement of left and right bialgebras	26
4	Splitting associative clusters into associative operations	27
4.1	n -Hypercubic $\epsilon[R]$ -bialgebras	27
4.2	Circular algebras	29
4.2.1	Constructions of n -circular algebras via right Baxter operators	29
4.2.2	Constructions of circular algebras	30
5	Octo-algebras	31
5.1	Three different ways to obtain quadri-algebras from an associative cluster	31
5.2	The augmented free octo-algebra and connected Hopf algebra	36
5.3	Construction of octo-algebras via three pairwise commuting Baxter operators	38
5.4	Another construction	40
6	Leibniz algebras and Poisson dialgebras from right Baxter operators	41
6.1	Theoretical constructions	41
6.2	Constructions from coassociative coverings of directed graphs	43
7	Conclusion and opennings	43

1 Introduction and notation

In the sequel k is a field of characteristic zero. Let (X, \bullet) be a k -algebra and $(\bullet_i)_{1 \leq i \leq N} : X^{\otimes 2} \rightarrow X$ be a family of binary operations on X . The notation $\bullet \longrightarrow \sum_i \bullet_i$ will mean $x \bullet y = \sum_i x \bullet_i y$, for all $x, y \in X$. We say that the operation \bullet *splits* into the N operations $\bullet_1, \dots, \bullet_N$, or that the operation \bullet is a *cluster of N (binary) operations*.

This paper is a continuation of [2, 9] on ϵ -bialgebras and commuting Baxter operators on the one hand and of [10, 12, 13] on weighted directed graphs on the other hand. A ϵ -bialgebra (A, μ, Δ) is an associative algebra (A, μ) together with a coassociative coalgebra (A, Δ) such that, for all $a, b \in A$, $\Delta(ab) := \Delta(a)b + a\Delta(b)$. Such bialgebras appeared for the first time in the work of Joni and Rota [7], see also [1, 2, 9, 22]. However, other structures, corresponding to the equality $\Delta(ab) := \Delta(a)b$ or $\Delta(ab) := a\Delta(b)$ can be met in algebraic descriptions of some substitutions in symbolic dynamics (see Section 1) or in the work of Khovanov [8] on 2-dimensional topological quantum field theory. They are called left or right ϵ -bialgebras and are defined as follows.

Definition 1.1 [$\epsilon[R]$ -, $\epsilon[L]$ -Bialgebra] A $\epsilon[R]$ -bialgebra (resp. $\epsilon[L]$ -bialgebra) (A, μ, Δ) is an associative algebra (A, μ) together with a coassociative coalgebra (A, Δ) such that, for all $a, b \in A$, $\Delta(ab) := a\Delta(b)$, (resp. $\Delta(ab) := \Delta(a)b$). Similarly, a $\epsilon'[R]$ -bialgebra (resp. $\epsilon'[L]$ -bialgebra) (A, μ, Δ) is an associative algebra (A, μ) together with a coalgebra (A, Δ) not supposed to be coassociative such that, for all $a, b \in A$, $\Delta(ab) := a\Delta(b)$, (resp. $\Delta(ab) := \Delta(a)b$). The linear map $\Delta : A \rightarrow A^{\otimes 2}$ is said to be a *co-operation*.

To any ϵ -bialgebras, commuting pairs of Baxter operators can be introduced [2]. Let (A, μ) be an associative algebra. Recall that a Baxter operator is a linear map $\zeta : A \rightarrow A$ such that, for all $x, y \in A$,

$$\zeta(x)\zeta(y) = \zeta(\zeta(x)y + x\zeta(y)).$$

Baxter operators on associative algebras appear originally in a work of G. Baxter [3] and the importance of such a map was stressed by G.-C. Rota in [21]. See also [1, 2, 4, 9] for more details.

Similarly, the notion of right Baxter operators (resp. left Baxter operators), i.e., linear maps $\zeta : A \rightarrow A$ verifying $\zeta(x)\zeta(y) = \zeta(\zeta(x)y)$ (resp. $\zeta(x)\zeta(y) = \zeta(x\zeta(y))$) on associative algebras can be introduced. Such operators appear in the study of $\epsilon'[R]$ -bialgebras and their left versions². The novelty lies in $\epsilon'[R]$ -bialgebras where co-operations are not supposed to be coassociative since right Baxter operators can be introduced as well as associative structures, called *L-anti-dipterous algebras* on $\text{End}(A)$, the k -algebra of linear maps from A to A . The case of $\epsilon[R]$ -bialgebras, where co-operations are supposed to be coassociative, produce left Baxter operators which commute with previous right ones. Other types of associative algebras, presenting symmetries, will appear. This is studied in Section 2. Examples are given. We also recall the definition of quadri-algebras [2] and give other examples of ϵ -bialgebras. In an example already described in [9], we relate ϵ -bialgebra structures to a toy-model of virus attacks inside a host cell.

To describe some mathematical objects, such as weighted directed graphs [13] or how graphs can be covered by coassociative structures, [10, 12], is possible only with the help of several co-operations. That is why in Section 3 and 4, we explore what occurs if associative algebras have two co-operations, related by an entanglement condition, which well behave with the underlying product. Such descriptions also produce commuting Baxter operators. Many cases are undertaken. This leads to the constructions of n -hypercubic algebras, n -circular algebras, both related to the notion of Hochschild 2-cocycles.

Up to now, we have worked with two commuting Baxter operators and their right or left versions. What happens if three Baxter operators commute pairwise? In section 5, we introduce particular k -vector spaces with two coproducts, called *L-circular ϵ -bialgebras*. Via such structures, three pairwise commuting Baxter operators are created. In the case of ϵ -bialgebras, two commuting Baxter operators were created giving birth to quadri-algebras [2], i.e., k -vector spaces Q equipped with four binary operations $\searrow, \nearrow, \swarrow, \nwarrow : Q^{\otimes 2} \rightarrow Q$ and obeying 9 compatibility axioms. One of the interests of such algebras is to produce an associative algebra (Q, \star) such that \star is the following associative cluster $\star \rightarrow \searrow + \nearrow + \swarrow + \nwarrow$. The second and main interest is to produce two dendriform dialgebras by combining these four operations, the

²In the sequel of this introduction, we will focus mainly on right versions.

two dendriform dialgebras (Q, \prec, \succ) and (Q, \wedge, \vee) being such that $\star \rightarrow \prec + \succ$ and $\star \rightarrow \wedge + \vee$. In the case of three pairwise commuting Baxter operators, the relevant notion is called octo-algebras. Octo-algebras are associative algebras whose associative product is a cluster of eight operations. By combining them, three quadri-algebras (or six dendriform dialgebras) sharing the same associative product can be produced. Examples are given, one of them coming from discrete mathematics. This work ends with Section 6 by showing how to produce, from right Baxter operators, Leibniz and Poisson dialgebras, notion introduced in [16, 17]. Examples from coassociative coverings of some directed graphs are also given.

The background to make this article as self-contained as possible is now presented.

Definition 1.2 [Dendriform dialgebra [16]] A *dendriform dialgebra* is a k -vector space E equipped with two binary operations: $\prec, \succ: E^{\otimes 2} \rightarrow E$, satisfying the following relations for all $x, y \in E$:

$$(x \prec y) \prec z = x \prec (y \star z), \quad (x \succ y) \prec z = x \succ (y \prec z), \quad (x \star y) \succ z = x \succ (y \succ z),$$

where by definition $x \star y := x \prec y + x \succ y$, for all $x, y \in E$. The dendriform dialgebra (E, \star) is then an associative algebra such that $\star \rightarrow \prec + \succ$. Observe that these axioms are globally invariant under the transformation $x \prec^{op} y := y \succ x$, $x \succ^{op} y := y \prec x$. A dendriform dialgebra is said to be *commutative* if $x \prec y := y \succ x$. Such algebras are also called *Zinbiel algebras* [16]. The free dendriform dialgebra is related to planar (rooted) binary trees.

Definition 1.3 [L-Algebra] A *L-algebra* $(A, \succ \dashrightarrow \prec)$ is a k -vector space equipped with two binary operations, $\succ, \prec: A^{\otimes 2} \rightarrow A$ *linked* for all $x, y, z \in A$ by,

$$(L.) \quad (x \succ y) \prec z = x \succ (y \prec z) \equiv \succ \dashrightarrow \prec.$$

A *L-algebra* is said to be *associative* if both \prec and \succ are associative. The free *L-algebra* is related to planar (rooted) binary trees [19]. The importance of *L-coalgebras* can be found in [13] and in the notion of coverings or tilings of directed graphs by coassociative structures [10, 12]. By the notation $\prec \dashrightarrow \succ$, we mean $(x \prec y) \succ z = x \prec (y \succ z)$. Similarly, for the conversion, $\Delta_{\prec} \dashrightarrow \Delta_{\succ}$ means $(\Delta_{\prec} \otimes id)\Delta_{\succ} := (id \otimes \Delta_{\succ})\Delta_{\prec}$. Such an equality will be called an *entanglement*.

Definition 1.4 [L-Dipterous algebra] A *dipterous algebra* (A, \star, \succ) is an associative algebra (A, \star) equipped with a left module on itself, i.e., for all $x, y, z \in A$,

$$(As.) \quad (x \star y) \star z = x \star (y \star z), \quad (Dip.) \quad (x \star y) \succ z = x \succ (y \succ z).$$

The free dipterous algebra is related to two copies of planar rooted trees [18]. Similarly, a *L-dipterous algebra* (A, \star, \succ) is a dipterous algebra verifying an extra condition:

$$(L.) \quad (x \succ y) \star z = x \succ (y \star z),$$

for all $x, y \in A$. A *L-anti-dipterous algebra* is a k -vector space A equipped with two binary operations $\star, \prec: A^{\otimes 2} \rightarrow A$, obeying,

$$(As.) \quad (x \star y) \star z = x \star (y \star z), \quad (A.Dip.) \quad (x \prec y) \prec z = x \prec (y \star z), \quad (L.) \quad (x \star y) \prec z = x \star (y \prec z).$$

In terms of category, we get,

$$\text{L-Dip.} \longrightarrow \text{Dip.} \longrightarrow \text{As.} \quad \text{and} \quad \text{L-A.Dip.} \longrightarrow \text{A.Dip.} \longrightarrow \text{As.}$$

All these k -vector spaces are said to be algebras of a certain type P . To better formulate this sentence, we need the operadic language, see for instance [5, 6, 14]. Let us briefly expose what this is. Let P be a type of algebras and $P(V)$ be the free P -algebra on the k -vector space V , i.e., by definition, a k -algebra of type P equipped with a map $i : V \mapsto P(V)$ which satisfies the following universal property: for any linear map $f : V \rightarrow A$, where A is a k -algebra of type P , there exists a unique morphism of algebra of type P , $\bar{f} : P(V) \rightarrow A$ such that $\bar{f} \circ i = f$. Suppose $P(V) := \bigoplus_{n \geq 1} P(n) \otimes_{S_n} V^{\otimes n}$, where $P(n)$ are right S_n -modules. We view P as an endofunctor on the category of k -vector spaces. The structure of the free P -algebra of $P(V)$ induces a natural transformation $\pi : P \circ P \rightarrow P$ as well as $u : Id \rightarrow P$ verifying usual associativity and unitarity axioms. An *algebraic operad* is then a triple (P, π, u) . A *P -algebra* or algebra of type P is then a k -vector space A together with a linear map $\pi_A : P(A) \rightarrow A$ such that $\pi_A \circ \pi(A) = \pi_A \circ P(\pi_A)$ and $\pi_A \circ u(A) = Id_A$. The k -vector space $P(n)$ is the space of n -ary operations for P -algebras. In the sequel we suppose there is, up to homotheties, a unique 1-ary operation, the identity, i.e., $P(1) := k.Id$ and that all possible operations are generated by composition from $P(2)$, i.e., by binary operations. The operad is said to be *binary*. It is said to be *quadratic* if all the relations between operations are consequences of relations described exclusively with the help of monomials with two operations. An operad is said to be *non-symmetric* if, in the relations, the variables x, y, z appear in the same order. In this case, the k -vector space $P(n)$ can be written like $P(n) := P'(n) \otimes k[S_n]$, where $P'(n)$ is also a k -vector space. The free P -algebra is then entirely induced by the free P -algebra on one generator $P(k) := \bigoplus_{n \geq 1} P'(n)$. The generating function of the operad P is given by:

$$f^P(x) := \sum (-1)^n \frac{\dim P(n)}{n!} x^n := \sum (-1)^n \dim P'(n) x^n.$$

In this paper, operads will be binary, quadratic and non-symmetric. Their sequences $(\dim P'(n))_{n \geq 1}$, at least their starts will be indicated.

Definition 1.5 [Baxter operators] Let (P, π, u) be a binary, quadratic and non-symmetric operad. A *Baxter operator*, (resp. *right Baxter operator*), (resp. *left Baxter operator*) on the k -algebra A of type P , is a linear map $\zeta : A \rightarrow A$, such that, for all $x, y \in A$ and for all $\circ \in P(2)$,

$$\zeta(x) \circ \zeta(y) = \zeta(\zeta(x) \circ y + x \circ \zeta(y)),$$

(resp. $\zeta(x) \circ \zeta(y) = \zeta(\zeta(x) \circ y)$), (resp. $\zeta(x) \circ \zeta(y) = \zeta(x \circ \zeta(y))$).

If (A, μ) is an associative k -algebra, then the following operations $a(x \otimes y) := ax \otimes y$ and $(x \otimes y)a := x \otimes ya$ defined for all $a, x, y \in A$ turn $A^{\otimes 2}$ into a A -bimodule. The k -vector space $\text{End}(A)$ of linear endomorphisms of A will be viewed as an associative algebra under composition denoted simply by concatenation TS , for $T, S \in \text{End}(A)$. The existence of a co-operation $\Delta : A \rightarrow A^{\otimes 2}$ endows $\text{End}(A)$ with the so-called *convolution product* $*$, defined by $T * S := \mu(T \otimes S)\Delta$, for all $T, S \in \text{End}(A)$. It is associative if the co-operation Δ is coassociative.

2 Algebraic structures related to one co-operation

2.1 $\epsilon'[R]$ -Bialgebras and L -anti-dipterous algebras

In this subsection, we show that associative structures can pop up from non-coassociative co-operations provided they well behave with the underlying associative product.

Lemma 2.1 *Let (A, μ) be an associative algebra and $\zeta : A \rightarrow A$ be a right Baxter operator. Define the binary operations, $\overleftarrow{\star}_\zeta, \prec_\zeta : A^{\otimes 2} \rightarrow A$ by:*

$$x \overleftarrow{\star}_\zeta y := \zeta(x)y, \quad x \prec_\zeta y := x\zeta(y), \quad \forall x, y \in A.$$

Then, $(A, \overleftarrow{\star}_\zeta, \prec_\zeta)$ is a L -anti-dipterous algebra.

Proof: Let (A, μ) be an associative algebra and $\zeta : A \rightarrow A$ be a right Baxter operator, i.e., for all $x, y \in A$,

$$\zeta(x)\zeta(y) := \zeta(\zeta(x)y).$$

Let us show for instance that the binary operation $\overleftarrow{\star}_\zeta$ is associative. Indeed, for all $x, y, z \in A$, $x \overleftarrow{\star}_\zeta (y \overleftarrow{\star}_\zeta z) := \zeta(x)\zeta(y)z$ and $(x \overleftarrow{\star}_\zeta y) \overleftarrow{\star}_\zeta z := \zeta(\zeta(x)y)z$. Checking the two other axioms of Definition 1.4 entails that $(A, \overleftarrow{\star}_\zeta, \prec_\zeta)$ is a L -anti-dipterous algebra. \square

One of the interest of L -anti-dipterous algebras can be found in the following theorem and its applications.

Theorem 2.2 *Let (A, μ, Δ) be a $\epsilon'[R]$ -bialgebra. Equip the algebra $\text{End}(A)$ with the convolution product $*$, not supposed to be associative here. Then, there exists a right Baxter operator, the right shift $\beta : \text{End}(A) \rightarrow \text{End}(A)$ given by $\beta(T) := id * T$, for all $T \in \text{End}(A)$. Set,*

$$T \overleftarrow{\star}_\beta S := \beta(T)S \quad \text{and} \quad T \prec_\beta S := T\beta(S), \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \overleftarrow{\star}_\beta, \prec_\beta)$ is a L -anti-dipterous algebra.

Proof: Let (A, μ, Δ) be a $\epsilon'[R]$ -bialgebra. For all $x \in A$, write, using the Sweedler notation, $\Delta(x) := x_{(1)} \otimes x_{(2)}$. Let $T, S \in \text{End}(A)$. On the one hand,

$$\begin{aligned} \beta(T)\beta(S)(x) &:= \beta(T)(x_{(1)}S(x_{(2)})), \\ &= \mu(id \otimes T)x_{(1)}S(x_{(2)})_{(1)} \otimes S(x_{(2)})_{(2)}, \\ &= x_{(1)}S(x_{(2)})_{(1)}T(S(x_{(2)})_{(2)}). \end{aligned}$$

On the other hand,

$$\begin{aligned} \beta(\beta(T)S)(x) &:= \mu(id \otimes \beta(T)S)(x_{(1)} \otimes x_{(2)}) = x_{(1)}\beta(T)(S(x_{(2)})), \\ &= x_{(1)}S(x_{(2)})_{(1)}T(S(x_{(2)})_{(2)}), \end{aligned}$$

proving that the right shift β is a right Baxter operator on $\text{End}(A)$. Applying Lemma 2.1 entails that $(\text{End}(A), \overleftarrow{\star}_\beta, \prec_\beta)$ is a L -anti-dipterous algebra. \square

Remark: From **non-coassociative** co-operations, associative algebras can be constructed via right Baxter operators. In terms of category, Theorem 2.2 yields,

$$\epsilon'[R] - \text{Bialg.} \longrightarrow \text{L-A.Dip.} \longrightarrow \text{As.}$$

Proposition 2.3 *Let $As(S)$ be the free associative k -algebra generated by a non-empty set S . Fix a co-operation $\Delta : kS \rightarrow kS^{\otimes 2}$ and extend it to a co-operation $\Delta_{\#} : As(S) \rightarrow As(S)^{\otimes 2}$ such that for all words say $s_1 \dots s_n$, $\Delta_{\#}(s_1 \dots s_n) := s_1 \dots s_{n-1} \Delta(s_n)$. Then, $(As(S), \Delta_{\#})$ is a $\epsilon'[R]$ -bialgebra.*

Proof: Straightforward by construction of the co-operation $\Delta_{\#}$. □

Example 2.4 [Weighted directed graphs and dynamics] Let $G = (G_0, G_1, s, t)$ be a directed graph, supposed to be locally-finite, row-finite, without sink and source, equipped with a family of weights $(w_v)_{v \in G_0}$ and such that $s \times t : G_1 \rightarrow G_0 \times G_0$ is injective. Consider the free k -vector space kG_0 spanned by G_0 . Identify any directed arrow $v \rightarrow w \in G_1$ with $v \otimes w$. The set G_1 is then viewed as a subset of $kG_0^{\otimes 2}$. The family of weights $(w_v)_{v \in G_0}$ is then viewed as a family of maps $w_v : F_v \rightarrow k$, where $F_v := \{a \in G_1, s(a) = v\}$. Define the co-operation $\Delta_M : kG_0 \rightarrow kG_0^{\otimes 2}$ as follows [13]:

$$\Delta_M(v) := \sum_{i: a_i \in F_v} w_v(a_i) v \otimes t(a_i),$$

for all $v \in G_0$. Extend it to $As(G_0)$ as in Proposition 2.3. Then, $(As(G_0), \Delta_M)_{\#}$ is a $\epsilon'[R]$ -bialgebra.

Example 2.5 [Substitutions and Language theory] Let S be a non-empty set. For all $s \in S$, suppose there exit substitutions of the form $s \mapsto s_1^i s_2^i$, $i \in I_s$, which appear with probability $\mathbb{P}(s \mapsto s_1^i s_2^i)$ and $\text{card}(I_s) < \infty$ for all $s \in S$. Consider the free k -vector space kS and define the co-operation $\Delta : kS \rightarrow kS^{\otimes 2}$ by,

$$\Delta(s) := \sum_{i \in I_s} \mathbb{P}(s \mapsto s_1^i s_2^i) s_1^i \otimes s_2^i.$$

Extend it to $As(S)$ as in Proposition 2.3. The k -vector space $(As(S), \Delta)_{\#}$ is then a $\epsilon'[R]$ -bialgebra.

All the possible dynamics of a string are then given by the operator $\mu\Delta$, where μ is the associative product representing the concatenation of two symbols. For instance, consider the start symbol at time $t = 0$, say s . Apply the operator Δ and μ to obtain,

$$\sum_{i \in I_s} \mathbb{P}(s \mapsto s_1^i s_2^i) s_1^i s_2^i,$$

which is all the possible strings at time $t = 1$. The probability to get the string, say $s_1^{i_0} s_2^{i_0}$, with $i_0 \in I_s$ at time $t = 1$, is then $\mathbb{P}(s \mapsto s_1^{i_0} s_2^{i_0})$ and so forth.

The dynamics of a string has another algebraic structure since from any $\epsilon'[R]$ -bialgebras, another L -anti-dipterous algebra can be constructed.

Proposition 2.6 *Let (A, μ, Δ) be a $\epsilon'[R]$ -bialgebra. Set $x \bowtie y := \mu(\Delta(x))y$ and $x \prec_A y := x\mu(\Delta(y))$ defined for all $x, y \in A$. Then, the k -vector space (A, \bowtie, \prec_A) is a L -anti-dipterous algebra.*

Proof: Straightforward by using $\mu(\Delta(xy)) = x\mu(\Delta(y))$, for all $x, y \in A$. \square

Proposition 2.6 gives another functor, $\epsilon'[R] - \text{Bialg.} \longrightarrow L\text{-A.Dip.}$

Remark: Denote by AL , the operad associated with L -anti-dipterous algebras. Then, the sequence $(AL(n))_{n \geq 1}$ starts with 1, 2, 5, 14, ...

2.2 On $\epsilon[R]$ -bialgebras

We suppose the co-operation Δ to be coassociative and study what kind of structures emerge.

Lemma 2.7 *Let (A, μ) be an associative algebra and $\theta : A \rightarrow A$ be a left Baxter operator. Define the binary operations $\overrightarrow{\star}_\theta, \succ_\theta : A^{\otimes 2} \rightarrow A$, by:*

$$x \overrightarrow{\star}_\theta y := x\theta(y), \quad x \succ_\theta y := \theta(x)y, \quad \forall x, y \in A.$$

Then, $(A, \overrightarrow{\star}_\theta, \succ_\theta)$ is a L -dipterous algebra.

Proof: Straightforward, since by definition left Baxter operators θ on (A, μ) obey the identity $\theta(x)\theta(y) := \theta(x\theta(y))$, for all $x, y \in A$. \square

As $\epsilon[R]$ -bialgebras are also $\epsilon'[R]$ -bialgebras, the operations $\overleftarrow{\star}_\beta$ and \prec_β defined in the previous subsection are related to $\overrightarrow{\star}_\gamma$ and \succ_γ via the following lemma.

Lemma 2.8 *Let (A, μ) be an associative algebra and $\zeta, (\theta) : A \rightarrow A$ be right (left) Baxter operators. Then,*

$$\begin{aligned} (x \overleftarrow{\star}_\zeta y) \overrightarrow{\star}_\theta z &= x \overleftarrow{\star}_\zeta (y \overrightarrow{\star}_\theta z), \\ (x \succ_\theta y) \prec_\zeta z &= x \succ_\theta (y \prec_\zeta z), \end{aligned}$$

for all $x, y \in A$.

Proof: Straightforward. \square

Theorem 2.9 *Let (A, μ, Δ) be a $\epsilon[R]$ -bialgebra. Then, there exists a left Baxter operator, the left shift $\gamma : \text{End}(A) \rightarrow \text{End}(A)$ given by $T \mapsto T * \text{id}$. Set,*

$$T \overrightarrow{\star}_\gamma S := T\gamma(S), \quad \text{and} \quad T \succ_\gamma S := \gamma(T)S, \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \overrightarrow{\star}_\gamma, \succ_\beta)$, is a L -dipterous algebra.

Moreover, the operations described in Theorem 2.2 are related to the new operations by the following relations:

$$\begin{aligned} (R \overleftarrow{\star}_\beta S) \overrightarrow{\star}_\gamma T &= R \overleftarrow{\star}_\beta (S \overrightarrow{\star}_\gamma T), \\ (R \succ_\gamma S) \prec_\beta T &= R \succ_\gamma (S \prec_\beta T), \end{aligned}$$

for all $R, S, T \in \text{End}(A)$.

Proof: Let (A, μ, Δ) be a $\epsilon[R]$ -bialgebra. Equip $\text{End}(A)$ with the convolution product $*$. Fix $T, S \in \text{End}(A)$, $x \in A$ and write $\Delta(x) := x_{(1)} \otimes x_{(2)}$. On the one hand,

$$\begin{aligned} \gamma(T)\gamma(S)(x) &:= \gamma(T)(S(x_{(1)})x_{(2)}), \\ &= \mu(T \otimes id) S(x_{(1)})(x_{(2)})_{(1)} \otimes (x_{(2)})_{(2)}, \\ &= T(S(x_{(1)})(x_{(2)})_{(1)})(x_{(2)})_{(2)}. \end{aligned}$$

One the other hand,

$$\begin{aligned} \gamma(T\gamma(S))(x) &:= \mu(T\gamma(S) \otimes id)(x_{(1)} \otimes x_{(2)}), \\ &= T\gamma(S)(x_{(1)})x_{(2)}, \\ &= (T\mu(S \otimes id)(x_{(1)})_{(1)} \otimes (x_{(1)})_{(2)})x_{(2)}, \\ &= T(S((x_{(1)})_{(1)})(x_{(1)})_{(2)})x_{(2)}. \end{aligned}$$

Therefore,

$$\gamma(T)\gamma(S) = \gamma(T\gamma(S)),$$

holds since the coproduct Δ is coassociative. The linear map γ is thus a left Baxter operator. The other claims follow from Lemmas 2.7 and 2.8. \square

In terms of category, Theorem 2.9 gives,

$$\epsilon[R] - \text{Bialg.} \longrightarrow \text{L-Dip.} \longrightarrow \text{Dip.} \longrightarrow \text{As.}$$

The coassociativity of the coproduct Δ will imply that the operators β and γ will commute. Let us show what kind of structures appear.

Theorem 2.10 *Let (A, μ) be an associative algebra and $\zeta, (\theta) : A \rightarrow A$ be right (left) Baxter operators which commute, i.e., $\zeta\theta = \theta\zeta$. Then, the left Baxter operator θ is a left Baxter operator on the L -anti-dipterous algebra $(A, \overleftarrow{\star}_\zeta, \prec_\zeta)$, i.e., for all $x, y \in A$,*

$$\theta(x) \overleftarrow{\star}_\zeta \theta(y) = \theta(x \overleftarrow{\star}_\zeta \theta(y)), \quad \theta(x) \prec_\zeta \theta(y) = \theta(x \prec_\zeta \theta(y)).$$

Moreover, this entails the existence of four extra binary operations $\bullet_1, \bullet_2, \bullet_3, \bullet_4 : A^{\otimes 2} \rightarrow A$ defined by (EQ. Th. 2.10),

$$x \bullet_1 y := x \overleftarrow{\star}_\zeta \theta(y) := \zeta(x)\theta(y), \quad x \bullet_2 y := \theta(x) \overleftarrow{\star}_\zeta y := \theta\zeta(x)y,$$

$$x \bullet_3 y := \theta(x) \prec_\zeta y := \theta(x)\zeta(y), \quad x \bullet_4 y := x \prec_\zeta \theta(y) := x\theta\zeta(y),$$

verifying the matrix of relations $([M]_{ij})_{(i,j=1,\dots,3)}$:

$$(x \bullet_4 y) \bullet_4 z = x \bullet_4 (y \bullet_1 z); \quad (x \bullet_3 y) \bullet_4 z = x \bullet_3 (y \bullet_1 z); \quad (x \bullet_1 y) \bullet_4 z = x \bullet_1 (y \bullet_4 z);$$

$$(x \bullet_4 y) \bullet_3 z = x \bullet_3 (y \bullet_2 z); \quad (x \bullet_2 y) \bullet_4 z = x \bullet_2 (y \bullet_4 z); \quad (x \bullet_1 y) \bullet_1 z = x \bullet_1 (y \bullet_1 z);$$

$$(x \bullet_1 y) \bullet_2 z = x \bullet_2 (y \bullet_2 z); \quad (x \bullet_1 y) \bullet_3 z = x \bullet_2 (y \bullet_3 z); \quad (x \bullet_2 y) \bullet_1 z = x \bullet_2 (y \bullet_1 z),$$

for all $x, y, z \in A$. We shall say that $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ has a $([M]_{ij})_{(i,j=1,\dots,3)}$ -algebra structure if its matrix of relations verifies $([M]_{ij})_{(i,j=1,\dots,3)}$, up to a permutation of S_9 .

Proof: Let (A, μ) be an associative algebra and $\zeta, (\theta) : A \rightarrow A$ be right (left) Baxter operators such that $\zeta\theta = \theta\zeta$. Fix $x, y, z \in A$. We have,

$$\begin{aligned} \theta(x) \overleftarrow{\star}_\zeta \theta(y) &= \zeta(\theta(x))\theta(y), \\ &= \theta(\zeta(x))\theta(y), \\ &= \theta(\zeta(x)\theta(y)), \\ &= \theta(x \overleftarrow{\star}_\zeta \theta(y)). \end{aligned}$$

Proceeding similarly for the operation \prec_ζ shows that θ is a left Baxter operator on $(A, \overleftarrow{\star}_\zeta, \prec_\zeta)$.

Let us check for instance $[M]_{11}$, i.e., the anti-dipterous relation: $(x \bullet_4 y) \bullet_4 z = x \bullet_4 (y \bullet_1 z)$.

$$\begin{aligned} (x \bullet_4 y) \bullet_4 z &:= (x \prec_\zeta \theta(y)) \prec_\zeta \theta(z), \\ &:= x \prec_\zeta (\theta(y) \overleftarrow{\star}_\zeta \theta(z)), \\ &:= x \prec_\zeta \theta(y \overleftarrow{\star}_\zeta \theta(z)), \\ &:= x \bullet_4 (y \bullet_1 z). \end{aligned}$$

Proceeding similarly for the 8 other equations shows that $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ has a $([M]_{ij})_{(i,j)}$ -algebra structure. \square

Keep notation of Theorem 2.10 and observe that $(A, \bullet_1, \bullet_4)$ is a L -anti-dipterous algebra whereas $(A, \bullet_1, \bullet_2)$ is a L -dipterous algebra. Therefore, in terms of category, Theorem 2.10 gives the following commutative diagram.

$$\begin{array}{ccccc} & & & \text{L-Dip.} & \\ & \nearrow & & \searrow & \\ \epsilon[R] - \text{Bialg.} & \longrightarrow & ([M]_{(ij)}) & \longrightarrow & \text{As.} \\ & \searrow & & \nearrow & \\ & & & \text{L-A.Dip.} & \end{array}$$

Remark: In Theorem 2.10 the rôles play by ζ and θ can be inversed, $\overleftarrow{\star}_\zeta$ becoming $\overrightarrow{\star}_\theta$ and \prec_ζ becoming \succ_θ . Hence, ζ becomes a right Baxter operator on the L -dipterous algebra $(A, \overrightarrow{\star}_\theta, \succ_\theta)$. In this case we will obtain new four binary operations, $\tilde{\bullet}_1, \tilde{\bullet}_2, \tilde{\bullet}_3, \tilde{\bullet}_4 : A^{\otimes 2} \rightarrow A$ defined by,

$$x \tilde{\bullet}_1 y := x \overrightarrow{\star}_\theta \zeta(y), \quad x \tilde{\bullet}_2 y := \zeta(x) \overrightarrow{\star}_\theta y,$$

$$x \tilde{\bullet}_3 y := \zeta(x) \succ_{\theta} y, \quad x \tilde{\bullet}_4 y := x \succ_{\theta} \zeta(y).$$

However, observe that the new matrix of relations can be obtained from (EQ. Th. 2.10) by the permutation σ defined by $\sigma(1) = 2$, $\sigma(2) = 3$, $\sigma(3) = 4$, $\sigma(4) = 1$. For instance, $[M]_{11}$ becomes,

$$[(x \tilde{\bullet}_{\sigma(4)} y) \tilde{\bullet}_{\sigma(4)} z = x \tilde{\bullet}_{\sigma(4)} (y \tilde{\bullet}_{\sigma(1)} z)] := [(x \tilde{\bullet}_1 y) \tilde{\bullet}_1 z = x \tilde{\bullet}_1 (y \tilde{\bullet}_2 z)].$$

However,

$$\begin{aligned} (x \tilde{\bullet}_1 y) \tilde{\bullet}_1 z &:= (x \overrightarrow{\star}_{\theta} \xi(y)) \overrightarrow{\star}_{\theta} \xi(z) = x \overrightarrow{\star}_{\theta} (\xi(y) \overrightarrow{\star}_{\theta} \xi(z)), \\ &= x \overrightarrow{\star}_{\theta} \xi((\xi(y) \overrightarrow{\star}_{\theta} z)), \\ &= x \tilde{\bullet}_1 (y \tilde{\bullet}_2 z). \end{aligned}$$

Proposition 2.11 [Opposite structure of $([M]_{ij})_{(i,j)}$ -algebras] Let $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ be a $([M]_{ij})_{(i,j)}$ -algebra. Define new binary operations $\bullet_1^{op}, \bullet_2^{op}, \bullet_3^{op}, \bullet_4^{op} : A^{\otimes 2} \rightarrow A$ by,

$$x \bullet_1^{op} y := y \bullet_1 x, \quad x \bullet_3^{op} y := y \bullet_3 x, \quad x \bullet_2^{op} y := y \bullet_4 x, \quad x \bullet_4^{op} y := y \bullet_2 x,$$

for all $x, y \in A$. Then, $(A, \bullet_1^{op}, \bullet_2^{op}, \bullet_3^{op}, \bullet_4^{op})$ is still a $([M]_{ij})_{(i,j=1,\dots,3)}$ -algebra.

Proof: Consider the matrix $([M]_{ij})_{(i,j)}$ stated in Theorem 2.10. Denote by $([M]_{ij}^{op})_{(i,j)}$ the matrix obtained from $([M]_{ij})_{(i,j)}$ by replacing $[M]_{ij}$ by the transformation stated in Proposition 2.11. Observe the existence of an axial symmetry around the row 2. Indeed, for all $j = 1, \dots, 3$ $[M]_{2j} := [M]_{2j}^{op}$, $[M]_{1j} := [M]_{3j}^{op}$ and $[M]_{3j} := [M]_{1j}^{op}$. \square

A $([M]_{ij})_{(i,j)}$ -algebra $(A, \bullet_1^{op}, \bullet_2^{op}, \bullet_3^{op}, \bullet_4^{op})$ is said to be *commutative* when it coincides with its opposite structure, i.e., $x \bullet_1 y = y \bullet_1 x$, $x \bullet_2 y = y \bullet_4 x$ and $x \bullet_3 y = y \bullet_3 x$.

Remark: The operad $P_{([M]_{ij})_{(i,j)}}$ associated with algebras of type $([M]_{ij})_{(i,j)}$ is binary, quadratic and non-symmetric. The sequence $(\dim P_{([M]_{ij})_{(i,j)}}(n))_{n \geq 1}$ starts with 1, 4, 23, 160, ...

Corollary 2.12 Let (A, μ, Δ) be a $\epsilon[R]$ -bialgebra. Then, there exists a structure of $([M]_{ij})_{(i,j)}$ -algebra on $\text{End}(A)$.

Proof: Keep notation of Theorem 2.10 and apply it to the right Baxter operator β which commutes with the left Baxter operator γ , where $\beta(T) := id * T$ and $\gamma(T) := T * id$, for all $T \in \text{End}(A)$. \square

Corollary 2.13 Let (A, μ) be an associative algebra and $\zeta, (\theta) : A \rightarrow A$ be two right Baxter operators which commute, i.e., $\zeta\theta = \theta\zeta$. Then, the right Baxter operator θ is a right Baxter operator on the L -anti-dipterous algebra $(A, \overleftarrow{\star}_{\zeta}, \prec_{\zeta})$, i.e., for all $x, y \in A$,

$$\theta(\theta(x) \overleftarrow{\star}_{\zeta} y) = \theta(\theta(x) \overleftarrow{\star}_{\zeta} y), \quad \theta(x) \prec_{\zeta} \theta(y) = \theta(\theta(x) \prec_{\zeta} y).$$

Moreover, this entails the existence of four extra binary operations $\bullet_1, \bullet_2, \bullet_3, \bullet_4 : A^{\otimes 2} \rightarrow A$, defined in Theorem 2.10, and verifying the matrix $([M_1]_{ij})_{(i,j=1,\dots,3)}$,

$$\begin{aligned} (x \bullet_2 y) \bullet_2 z &= x \bullet_2 (y \bullet_2 z); & (x \bullet_1 y) \bullet_4 z &= x \bullet_1 (y \bullet_3 z); & (x \bullet_2 y) \bullet_1 z &= x \bullet_2 (y \bullet_1 z); \\ (x \bullet_3 y) \bullet_4 z &= x \bullet_3 (y \bullet_1 z); & (x \bullet_2 y) \bullet_4 z &= x \bullet_2 (y \bullet_4 z); & (x \bullet_1 y) \bullet_1 z &= x \bullet_1 (y \bullet_2 z); \\ (x \bullet_2 y) \bullet_3 z &= x \bullet_2 (y \bullet_3 z); & (x \bullet_3 y) \bullet_3 z &= x \bullet_3 (y \bullet_2 z); & (x \bullet_4 y) \bullet_4 z &= x \bullet_4 (y \bullet_2 z). \end{aligned}$$

We shall say that the k -vector space $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ is an algebra of type $([M_1]_{ij})_{(i,j=1,\dots,3)}$ if its matrix of relations is $([M_1]_{ij})_{(i,j=1,\dots,3)}$ up to a permutation of S_9 .

Proof: As the Baxter operator θ is no longer left but right, this implies the existence of the following relations,

$$\begin{aligned} x \bullet_1 (y \bullet_i z) &\longleftrightarrow x \bullet_1 (y \bullet_j z), & x \bullet_4 (y \bullet_i z) &\longleftrightarrow x \bullet_4 (y \bullet_j z), \\ (x \bullet_i y) \bullet_3 z &\longleftrightarrow (x \bullet_j y) \bullet_3 z, & (x \bullet_i y) \bullet_2 z &\longleftrightarrow (x \bullet_j y) \bullet_2 z, \end{aligned}$$

with if $i = 1$, (resp. $i = 4$) then j has to be removed by 2 (resp. by 3). Apply these relations to the matrix $([M]_{ij})_{(i,j=1,\dots,3)}$ of Theorem 2.10 to obtain the matrix $([M_1]_{ij})_{(i,j=1,\dots,3)}$ up to a permutation of S_9 . \square

Keep notation of Corollary 2.13 and observe that both $(A, \bullet_2, \bullet_3)$ and $(A, \bullet_2, \bullet_1)$ are L -anti-pterous algebras. Therefore, Corollary 2.13 gives the following commutative diagram.

$$\begin{array}{ccc} & \text{L-A.Dip.} & \\ \nearrow & & \searrow \\ ([M_1]_{(ij)}) & \longrightarrow & \text{As.} \\ \searrow & & \nearrow \\ & \text{L-A.Dip.} & \end{array}$$

Remark: Denote by $P_{([M_1]_{ij})_{(i,j)}}$, the binary quadratic non-symmetric operad associated with algebra of type $([M_1]_{ij})_{(i,j=1,\dots,3)}$. The sequence $(P_{([M_1]_{ij})_{(i,j)}}(n))_{n \geq 1}$ starts with 1, 4, 23, 160, ...

Proposition 2.14 [Transpose of algebras of type $([M_1]_{ij})_{(i,j)}$] Let $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ be an algebra of type $([M_1]_{ij})_{(i,j)}$. Define for all $x, y \in A$ four binary operations, $\bullet_{1,t}, \bullet_{2,t}, \bullet_{3,t}, \bullet_{4,t} : A^{\otimes 2} \rightarrow A$ by,

$$(Tr\ 2.) \quad x \bullet_{1,t} y := x \bullet_3 y, \quad x \bullet_{3,t} y := x \bullet_1 y, \quad x \bullet_{2,t} y := x \bullet_2 y, \quad x \bullet_{4,t} y := x \bullet_4 y.$$

Denote by $([M_1]_{ij}^t)_{(i,j)}$ the matrix of relations obtained from $([M_1]_{ij})_{(i,j)}$ by applying the transformation $(Tr2.)$. Then, the k -vector space A equipped with these four binary operations is still an algebra of type $([M_1]_{ij})_{(i,j)}$.

Proof: Observe that the transformation $(Tr\ 2.)$ let the matrix $([M_1]_{ij})_{(i,j=1,\dots,3)}$ globally invariant since for all $i, j = 1, \dots, 3$, $[M_1]_{ij} = [M_1]_{ji}$. \square

Remark: Another extra algebraic structure can be exhibited. Let (A, μ, Δ) be a $\epsilon[R]$ -bialgebra. There exist 3 extras binary operations $\searrow, \nearrow, \nwarrow : A^{\otimes 2} \rightarrow A$, given by,

$$x \nearrow y := x \xleftarrow{\star} \beta \gamma(y) := \beta(x) \xrightarrow{\star} \gamma y, \quad x \nwarrow y := x \prec_{\beta} \gamma(y), \quad x \searrow y := \beta(x) \succ_{\gamma} y,$$

and verifying:

$$\begin{aligned} (x \nearrow y) \nearrow z &= x \nearrow (y \nearrow z), & (x \nwarrow y) \nwarrow z &= x \nwarrow (y \nearrow z), & (x \searrow y) \nearrow z &= x \searrow (y \nearrow z), \\ [(x \nearrow y) \nearrow z &= x \nearrow (y \nearrow z)], & (x \nearrow y) \searrow z &= x \searrow (y \searrow z), & (x \nearrow y) \nwarrow z &= x \nearrow (y \nwarrow z), \\ (x \searrow y) \nwarrow z &= x \searrow (y \nwarrow z). \end{aligned}$$

Remark: Observe that the first row describes L -anti-dipterous algebra axioms, the second one L -dipterous algebra axioms, these two structures being linked by the last relation. The sequence of dimensions of the associated operad starts with $1, 3, 12, \dots$

Let us construct examples of $\epsilon[R]$ -bialgebras.

Example 2.15 Fix $\lambda \in k$. Let A be the unital commutative k -algebra $k[X_1, \dots, X_n]/\mathcal{R}$, where \mathcal{R} is the set of relations $X_i X_j = 0$, for all $i = 1, \dots, n$. Set $\Sigma := \sum_{i=1, \dots, n} X_i$. Then, the following co-operation $\Delta_{\lambda} : A \rightarrow A^{\otimes 2}$, defined by $\Delta_{\lambda}(1) := 1 \otimes \Sigma + \Sigma \otimes 1 + \lambda \Sigma \otimes \Sigma$ and $\Delta_{\lambda}(X_i) := X_i \otimes \Sigma$ is coassociative and turns A into a $\epsilon[R]$ -bialgebra. There exists a linear map $\epsilon_{\lambda} : A \rightarrow k$ given by $\epsilon_{\lambda}(\Sigma) := 1$ and $\epsilon_{\lambda}(1) := -\lambda$ verifying $(id \otimes \epsilon_{\lambda})\Delta_{\lambda}(1) = 1 = (\epsilon_{\lambda} \otimes id)\Delta_{\lambda}(1)$ and $(id \otimes \epsilon_{\lambda})\Delta_{\lambda}(X_i) = X_i$. Observe that in the unital commutative k -algebra $k[X]$, the coproduct is cocommutative. Such a structure appears in [8].

Proposition 2.16 Let S be a non-empty set and $\Delta : kS \rightarrow kS^{\otimes 2}$ be a coassociative co-operation. Consider the free associative k -algebra $As(S)$ generated by S . Extend the coproduct Δ to $As(S)$ by considering $\Delta_{\#} : As(S) \rightarrow As(S)^{\otimes 2}$ defined by $\Delta_{\#}(s_1 \dots s_n) := s_1 \dots s_{n-1} \Delta(s_n)$, where $s_1, \dots, s_n \in S$. Then, $(As(S), \Delta_{\#})$ is a $\epsilon[R]$ -bialgebra. If $(\eta \otimes id)\Delta = id$ (resp. $(id \otimes \eta)\Delta = id$) holds for a linear map $\eta : kS \rightarrow k$ then so are the equations $(\eta_{\#} \otimes id)\Delta_{\#} = id$ (resp. $(id \otimes \eta_{\#})\Delta_{\#} = id$) for the map $\eta_{\#} : As(S) \rightarrow As(S) \oplus k$ given by $\eta_{\#}(s_1 \dots s_{n-1} s_n) := s_1 \dots s_{n-1} \eta(s_n)$, where $s_1, \dots, s_n \in S$.

Proof: Let S be a set and $\Delta : kS \rightarrow kS^{\otimes 2}$ be a coassociative co-operation. Consider the free associative k -algebra $As(S)$ generated by S . We have to show that $\Delta_{\#} : As(S) \rightarrow As(S)^{\otimes 2}$ is coassociative. Fix $s_1, \dots, s_n \in S$. As $s_1 \dots s_n \in As(S)$ and $\Delta(s_n) := (s_n)_{(1)} \otimes (s_n)_{(2)}$ we get:

$$(id \otimes \Delta_{\#})\Delta_{\#}(s_1 \dots s_n) := s_1 \dots s_{n-1} (s_n)_{(1)} \otimes \Delta((s_n)_{(2)}),$$

and

$$(\Delta_{\#} \otimes id)\Delta_{\#}(s_1 \dots s_n) := s_1 \dots s_{n-1} \Delta((s_n)_{(1)}) \otimes (s_n)_{(2)}.$$

Therefore, $(As(S), \Delta_{\#})$ is a $\epsilon[R]$ -bialgebra. □

Remark: As an application, we can deal with language theory and coassociative substitutions, see [11, 10]. Let S be a non-empty set. Suppose substitutions on S are of the form $s \mapsto s_1^i s_2^j$,

with $i \in I_s$ and $\text{card } I_s < \infty$. Consider the free k -vector space kS spanned by S and construct the co-operation $\Delta : s \mapsto \sum_{i \in I_s} s_1^i \otimes s_2^i$. If Δ is coassociative, then the set of substitutions on S is said to be *coassociative*. Such substitutions appear in an algebraic description of the chaotic map $x \mapsto 2x \bmod 1$, with $x \in [0, 1]$ and a quantisation of the Bernoulli walk [11]. They also appear in works on coassociative tilings and coverings of directed graphs [10, 12].

Example 2.17 [$M_2(k)$] We give now another method, inspired from [1] to obtain $\epsilon[R]$ -bialgebras.

Proposition 2.18 *Let (A, μ) be an associative algebra and J be a finite set. Fix $A_j, B_j \in A$, $j \in J$. Consider the co-operation $\Delta : A \rightarrow A^{\otimes 2}$ defined by $x \mapsto \Delta(x) := \sum_j x A_j \otimes B_j$. Then, (A, μ, Δ) is a $\epsilon[R]$ -bialgebra if,*

$$\sum_{i,j} A_i \otimes B_i A_j \otimes B_j = \sum_{i,j} A_i A_j \otimes B_j \otimes B_i.$$

Proof: Straightforward. □

Let (A, μ) be an associative algebra and $\lambda \in k$. Consider the co-operation $\Delta_1 : A \rightarrow A^{\otimes 2}$ defined by $x \mapsto \Delta_1(x) := x A_1 \otimes B_1$. Then the co-operation Δ_1 is coassociative if $A_1^2 = \lambda A_1$ and $B_1 A_1 = \lambda B_1$. For instance consider in $M_2(k)$ the following matrices:

$$A = \begin{pmatrix} \lambda & \lambda \\ 0 & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 0 & 0 \\ \lambda & \lambda \end{pmatrix}.$$

Equipped with this coproduct, $M_2(k)$ is a $\epsilon[R]$ -bialgebra.

2.3 $\epsilon'[L]$ -Bialgebras and L -anti-dipterous algebras

What was done with $\epsilon[R]$ -bialgebras can be realised with their left versions. The rôle played by the right Baxter operator β is now hold in the left version by the operator γ and conversely.

Theorem 2.19 *Let (A, μ, Δ) be a $\epsilon'[L]$ -bialgebra. Equip the k -algebra $\text{End}(A)$ with the convolution product $*$, not supposed to be associative here. Then, the left shift $\gamma : \text{End}(A) \rightarrow \text{End}(A)$ given by $T \mapsto T * \text{id}$ is a right Baxter operator. Set,*

$$T \overleftarrow{\star}_{\gamma} S := \gamma(T)S \quad \text{and} \quad T \prec_{\gamma} S := T\gamma(S), \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \overleftarrow{\star}_{\gamma}, \prec_{\gamma})$ is a L -anti-dipterous algebra.

Proof: Let (A, μ, Δ) be a $\epsilon'[L]$ -bialgebra. Fix $T, S \in \text{End}(A)$, $x \in A$ and set $\Delta(x) := x_{(1)} \otimes x_{(2)}$. On the one hand,

$$\begin{aligned} \gamma(T)\gamma(S)(x) &:= \gamma(T)(S(x_{(1)})x_{(2)}), \\ &= \mu(T \otimes \text{id})S(x_{(1)})_{(1)} \otimes S(x_{(1)})_{(2)}x_{(2)}, \\ &= T(S(x_{(1)})_{(1)})(S(x_{(1)})_{(2)}x_{(2)}). \end{aligned}$$

On the other hand,

$$\begin{aligned}\gamma(\gamma(T)S)(x) &:= \mu(\gamma(T)S \otimes id)(x_{(1)} \otimes x_{(2)}) = \gamma(T)S(x_{(1)})x_{(2)}, \\ &= T(S(x_{(1)})_{(1)})(S(x_{(1)})_{(2)})x_{(2)},\end{aligned}$$

proving that γ is a right Baxter operator. Applying Lemma 2.1 entails that $(\text{End}(A), \overleftarrow{\star}_\gamma, \prec_\gamma)$ is a L -anti-dipterous algebra. \square

In terms of category, we get:

$$\epsilon'[L] - \text{Bialg.} \longrightarrow \text{L-A.Dip.} \longrightarrow \text{As.}$$

Let us construct examples of $\epsilon'[L]$ -bialgebras.

Proposition 2.20 *Let $As(S)$ be the free associative k -algebra generated by a non-empty set S . Fix a co-operation $\Delta : kS \rightarrow kS^{\otimes 2}$ and extend it to a co-operation $\Delta_\# : As(S) \rightarrow As(S)^{\otimes 2}$ such that for all words say $s_1 \dots s_n$, $\Delta_\#(s_1 \dots s_n) := \Delta(s_1)s_2 \dots s_{n-1}s_n$. Then, $(As(S), \Delta_\#)$ is a $\epsilon'[L]$ -bialgebra.*

Proof: Straightforward by construction of the co-operation $\Delta_\#$. \square

Example 2.21 [Weighted directed graphs and dynamics] Keep notation of example 2.4. Define the co-operation $\tilde{\Delta}_M : kG_0 \rightarrow kG_0^{\otimes 2}$ as follows [13]:

$$\tilde{\Delta}_M(v) := \sum_{i: a_i \in P_v} w_{s(a_i)}(a_i) s(a_i) \otimes v,$$

where $P_v := \{a \in G_1, t(a) = v\}$, for all $v \in G_0$. Then, $(As(G_0), \tilde{\Delta}_{\#M})$ is a $\epsilon'[L]$ -bialgebra.

More generally, we get,

Example 2.22 [Substitutions and Language theory] Let S be a non-empty set. For all $s \in S$, suppose there exit substitutions of the form $s \mapsto s_1^i s_2^i$, $i \in I_s$ with $\text{card}(I_s) < \infty$, which appear with probability $\mathbb{P}(s \mapsto s_1^i s_2^i)$. Consider the free k -vector space kS spanned by S and define the co-operation $\Delta : kS \rightarrow kS^{\otimes 2}$ by,

$$\Delta(s) := \sum_{i \in I_s} \mathbb{P}(s \mapsto s_1^i s_2^i) s_1^i \otimes s_2^i.$$

Then, extend it to $As(S)$ as in Proposition 2.20. The k -vector space $(As(S), \Delta_\#)$ is a $\epsilon'[L]$ -bialgebra.

From any $\epsilon'[L]$ -bialgebras, a L -dipterous algebra can be constructed.

Proposition 2.23 *Let (A, μ, Δ) be a $\epsilon'[L]$ -bialgebra. Set $x \succ_A y := \mu(\Delta(x))y$ and $x \bowtie y := x\mu(\Delta(y))$ defined for all $x, y \in A$. Then, the k -vector space (A, \bowtie, \succ_A) is a L -dipterous algebra.*

Proof: Straightforward by using $\mu(\Delta(xy)) = \mu(\Delta(x))y$, for all $x, y \in A$. \square

Remark: Applied this proposition to model substitutions in symbolic dynamics via the associative product \bowtie or the operation \succ_A .

2.4 On $\epsilon[L]$ -bialgebras

In this subsection, the co-operation Δ is supposed to be coassociative.

Theorem 2.24 *Let (A, μ, Δ) be a $\epsilon[L]$ -bialgebra. Then, the right shift $\beta : \text{End}(A) \rightarrow \text{End}(A)$ given by $T \mapsto id * T$ is a left Baxter operator. Set,*

$$T \overrightarrow{\star}_{\beta} S := T\beta(S), \quad \text{and} \quad T \succ_{\beta} S := \beta(T)S, \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \overrightarrow{\star}_{\beta}, \succ_{\beta})$ is a L -dipterous algebra. Therefore there exists a functor such that,

$$\epsilon[L] - \text{Bialg.} \longrightarrow L - \text{Dip.}$$

These two operations are related to \prec_{γ} and $\overleftarrow{\star}_{\gamma}$ from Theorem 2.19 by,

$$(R \overleftarrow{\star}_{\gamma} S) \overrightarrow{\star}_{\beta} T = R \overleftarrow{\star}_{\gamma} (S \overrightarrow{\star}_{\beta} T),$$

$$(R \succ_{\beta} S) \prec_{\gamma} T = R \succ_{\beta} (S \prec_{\gamma} T),$$

for all $R, S, T \in \text{End}(A)$.

Proof: Let (A, μ, Δ) be a $\epsilon[L]$ -bialgebra. Equip $\text{End}(A)$ with the convolution product $*$. Fix $T, S \in \text{End}(A)$, $x \in A$ and set $\Delta(x) := x_{(1)} \otimes x_{(2)}$. On the one hand,

$$\begin{aligned} \beta(T)\beta(S)(x) &:= \beta(T)(x_{(1)}S(x_{(2)})), \\ &= \mu(id \otimes T)(x_{(1)})_{(1)} \otimes (x_{(1)})_{(2)}S(x_{(2)}), \\ &= (x_{(1)})_{(1)}T((x_{(1)})_{(2)}S(x_{(2)})). \end{aligned}$$

One the other hand,

$$\begin{aligned} \beta(T\beta(S))(x) &:= \mu(id \otimes T\beta(S))(x_{(1)} \otimes x_{(2)}), \\ &= x_{(1)}T\beta(S)(x_{(2)}), \\ &= x_{(1)}T\mu(id \otimes S)((x_{(2)})_{(1)} \otimes (x_{(2)})_{(2)}), \\ &= x_{(1)}T((x_{(2)})_{(1)}S((x_{(2)})_{(2)})). \end{aligned}$$

Since the coproduct Δ is coassociative,

$$\beta(T)\beta(S) = \beta(T\beta(S)),$$

holds. The linear map β is a left Baxter operator. The other claims follow from Lemmas 2.7 and 2.8. \square

Remark: In the left version, the rôles of the operators β and γ are reversed.

Corollary 2.25 *Let (A, μ, Δ) be a $\epsilon[L]$ -bialgebra. Then, there exists a $([M]_{ij})_{(i,j)}$ -algebra structure on $\text{End}(A)$.*

Proof: Keep notation of Theorem 2.10 and apply it to the right Baxter operator γ which commutes with the left Baxter operator β , where $\beta(T) := id * T$ and $\gamma(T) := T * id$, for all $T \in \text{End}(A)$. \square

In terms of category, the following diagram commutes.

$$\begin{array}{ccccc}
 & & \text{L-Dip.} & & \\
 & \nearrow & & \searrow & \\
 \epsilon[L] - \text{Bialg.} & \longrightarrow & ([M]_{ij}) & \longrightarrow & \text{As.} \\
 & \searrow & & \nearrow & \\
 & & \text{L-A.Dip.} & &
 \end{array}$$

Proposition 2.26 *Let (A, μ) be an associative algebra and $\zeta, \theta : A \rightarrow A$ be two left Baxter operators which commute, i.e., $\zeta\theta = \theta\zeta$. Then, the left Baxter operator θ is a left Baxter operator on the L -dipterous algebra $(A, \overrightarrow{\star}_\zeta, \succ_\zeta)$, i.e., for all $x, y \in A$,*

$$\theta(x) \overrightarrow{\star}_\zeta \theta(y) = \theta(x \overrightarrow{\star}_\zeta \theta(y)), \quad \theta(x) \succ_\zeta \theta(y) = \theta(x \succ_\zeta \theta(y)).$$

Moreover, this entails the existence of four extra binary operations $\bullet_1, \bullet_2, \bullet_3, \bullet_4 : A^{\otimes 2} \rightarrow A$, defined by,

$$\begin{aligned}
 x \bullet_4 y &:= x \overrightarrow{\star}_\zeta \theta(y) := x \zeta(\theta(y)), & x \bullet_3 y &:= \theta(x) \overrightarrow{\star}_\zeta y := \theta(x) \zeta(y), \\
 x \bullet_2 y &:= \theta(x) \succ_\zeta y := \zeta(\theta(x))y, & x \bullet_1 y &:= x \succ_\zeta \theta(y) := \zeta(x) \theta(y),
 \end{aligned}$$

verifying the matrix of relations $([M_2]_{ij})_{(i,j)=1,\dots,3}$:

$$\begin{aligned}
 (x \bullet_4 y) \bullet_4 z &= x \bullet_4 (y \bullet_4 z); & (x \bullet_3 y) \bullet_1 z &= x \bullet_2 (y \bullet_1 z); & (x \bullet_1 y) \bullet_4 z &= x \bullet_1 (y \bullet_4 z); \\
 (x \bullet_1 y) \bullet_3 z &= x \bullet_2 (y \bullet_3 z); & (x \bullet_2 y) \bullet_4 z &= x \bullet_2 (y \bullet_4 z); & (x \bullet_4 y) \bullet_1 z &= x \bullet_1 (y \bullet_1 z); \\
 (x \bullet_3 y) \bullet_4 z &= x \bullet_3 (y \bullet_4 z); & (x \bullet_4 y) \bullet_3 z &= x \bullet_3 (y \bullet_3 z); & (x \bullet_4 y) \bullet_2 z &= x \bullet_2 (y \bullet_2 z).
 \end{aligned}$$

We shall say that the k -vector space $(A, \bullet_1, \bullet_2, \bullet_3, \bullet_4)$ is an algebra of type $([M_2]_{ij})_{(i,j)}$ if its matrix of relations is equal to $([M_2]_{ij})_{(i,j)}$ up to a permutation of S_9 .

Proof: Straightforward. \square

Observe that both the k -vector spaces $(A, \bullet_4, \bullet_1)$ and $(A, \bullet_4, \bullet_3)$ are L -dipterous algebras. Therefore the following diagram,

$$\begin{array}{ccccc}
 & & \text{L-Dip.} & & \\
 & \nearrow & & \searrow & \\
 ([M_2]_{ij}) & \longrightarrow & & \longrightarrow & \text{As.} \\
 & \searrow & & \nearrow & \\
 & & \text{L-Dip.} & &
 \end{array}$$

commutes.

Proposition 2.27 [*Transpose of an algebra of type* $([M_2]_{ij})_{(i,j)}$] *Keep notation of Proposition 2.26. Let A be a $([M_2]_{ij})_{(i,j)}$ -algebra. Define four new binary operations $\bullet_{i,t}$, $i = 1, \dots, 4$ by,*

$$\bullet_{1,t} \mapsto \bullet_3, \quad \bullet_{2,t} \mapsto \bullet_2, \quad \bullet_{3,t} \mapsto \bullet_1, \quad \bullet_{4,t} \mapsto \bullet_4.$$

Then, $(A, \bullet_{1,t}, \bullet_{2,t}, \bullet_{3,t}, \bullet_{4,t})$ is still an algebra of type $([M_2]_{ij})_{(i,j)}$.

Proof: Denote by $([M_2]_{ij}^t)_{i,j}$, the matrix of relations obtained from $([M_2]_{ij})_{i,j}$ by applying the said transformation. Then, observe that $[M_2]_{ij} := [M_2]_{ji}^t$, for all $i, j = 1, \dots, 3$. \square

Remark: Keep notation of Proposition 2.26. The rôle of the operators γ and ξ can be exchanged. In this case, the $([M_2]_{ij})_{(i,j)}$ -algebra we obtain is the transpose of the first established in Proposition 2.26.

Proposition 2.28 *Let A be a $([M_2]_{ij})_{(i,j)}$ -algebra. Consider the following binary operations*

$$x \bullet_{1,op} y := y \bullet_1 x, \quad x \bullet_{2,op} y := y \bullet_4 x, \quad x \bullet_{3,op} y := y \bullet_3 x, \quad x \bullet_{4,op} y := y \bullet_2 x,$$

for all $x, y \in A$. Then, $(A, \bullet_{1,op}, \bullet_{2,op}, \bullet_{3,op}, \bullet_{4,op})$ is a $([M_1]_{ij})_{(i,j)}$ -algebra.

Proof: Denote by $([M_2]_{ij}^{op})_{(i,j)}$, the matrix of relations obtained from $([M_2]_{ij})_{(i,j)}$ by applying the said transformation. Observe that: $([M_2]_{ij}^{op}) = [M_1]_{ij}$, for all $i, j = 1, \dots, 3$, (and up to a permutation of S_9 if the matrices are written in a different order). \square

Proposition 2.28 can be re-written in terms of category. By observing that the transformation op is one-to-one, we get,

$$([M_2]_{ij}) \rightleftharpoons ([M_1]_{ij}).$$

2.5 ϵ -Bialgebras

An *infinitesimal bialgebra* (abbreviated ϵ -bialgebra) is a triple (A, μ, Δ) where (A, μ) is an associative algebra and (A, Δ) is a coassociative coalgebra such that for all $a, b \in A$, $\Delta(ab) = a_{(1)} \otimes a_{(2)}b + ab_{(1)} \otimes b_{(2)}$. Such bialgebras appeared for the first time in the work of Joni and Rota [7], see also [1, 2, 9, 22]. ϵ -Bialgebras allow the construction of quadri-algebras [2]. For completeness, some definitions and results are recalled.

Definition 2.29 [*Quadri-algebra*] A *quadri-algebra* Q is a k -vector space equipped with four binary operations $\nearrow, \searrow, \swarrow, \nwarrow: A^{\otimes 2} \rightarrow A$ obeying 9 relations. To state them, we introduce

the following sums.

$$\begin{aligned}
x \succ y &:= x \nearrow y + x \searrow y, \\
x \prec y &:= x \nwarrow y + x \swarrow y, \\
x \vee y &:= x \searrow y + x \swarrow y, \\
x \wedge y &:= x \nearrow y + x \nwarrow y,
\end{aligned}$$

and

$$x \bar{*} y := x \prec y + x \succ y = x \wedge y + x \vee y,$$

for all $x, y \in Q$. The compatibility axioms are presented in the following 3×3 matrix $([Q]_{ij})_{(i,j=1,\dots,3)}$.

$$\begin{aligned}
(x \nwarrow y) \nwarrow z &= x \nwarrow (y \bar{*} z); & (x \nearrow y) \nwarrow z &= x \nearrow (y \prec z); & (x \wedge y) \nearrow z &= x \nearrow (y \succ z); \\
(x \swarrow y) \nwarrow z &= x \swarrow (y \wedge z); & (x \searrow y) \nwarrow z &= x \searrow (y \nwarrow z); & (x \vee y) \nearrow z &= x \searrow (y \nearrow z); \\
(x \prec y) \swarrow z &= x \swarrow (y \vee z); & (x \succ y) \swarrow z &= x \searrow (y \swarrow z); & (x \bar{*} y) \searrow z &= x \searrow (y \searrow z).
\end{aligned}$$

Theorem 2.30 ([2]) *Let (A, μ, Δ) be a ϵ -bialgebra. Then, there exist two commuting Baxter operators β and γ and thus a quadri-algebra structure on $\text{End}(A)$ given by:*

$$T \swarrow S := \gamma(T)\beta(S), \quad T \nwarrow S := T\gamma\beta(S), \quad T \searrow S := \gamma\beta(T)S, \quad T \nearrow S := \beta(T)\gamma(S),$$

with $\beta(T) := id * T$ and $\gamma(T) := T * id$, for all $T, S \in \text{End}(A)$.

We recall the existence of a notion of commutative quadri-algebras and of transpose of quadri-algebras. The free quadri-algebra could be related to non-crossing connected graphs [2]. The augmented free quadri-algebra has a connected Hopf algebra structure [14]. All these terms are explained in Section 5. A quadri-algebra gives two dendriform dialgebras, the so-called vertical structure $Q_v := (Q, \wedge, \vee)$ and the so-called horizontal structure $Q_h := (Q, \prec, \succ)$. The associative product $\bar{*}$ is then a cluster of four operations $x \bar{*} y := x \nearrow y + x \searrow y + x \swarrow y + x \nwarrow y$. In terms of categories, the following diagram commutes:

$$\begin{array}{ccc}
\text{Quadri.} & \xrightarrow{F_v} & \text{DiDend.} \\
F_h \downarrow & \searrow f & \downarrow F \\
\text{DiDend.} & \xrightarrow{F} & \text{As.}
\end{array}$$

This diagram is commutative, i.e., $FF_v = f = FF_h$, where F_v (resp. F_h) is the functor giving the vertical structure (resp. the horizontal structure) of a quadri-algebra. Section 5 is devoted to a generalisation of this structure obtained by considering three commuting Baxter operators.

2.6 Construction of ϵ -bialgebras from weighted directed graphs

Let X, V be two non-empty sets. Let $As(V)$ be the free associative algebra generated by V . By $As(V)\langle X \rangle$, we mean the non-commutative algebra of polynomials with indeterminates belong-

ing to X with coefficients in $As(V)$, i.e., $x \in As(V)\langle X \rangle$ is a linear combinaison of the form $v_1 v_2 X_1 v_3 \dots v_p X_2 \dots$, with $X_i \in X$ and $v_j \in As(V)$.

Proposition 2.31 *Let X, V be two non-empty sets. Let $As(V)$ be the free associative algebra generated by V . Suppose there exists a coassociative co-operation $\Delta : kX \rightarrow kX^{\otimes 2}$. Then, the linear map $\Delta_{\#} : As(V)\langle X \rangle \rightarrow As(V)\langle X \rangle^{\otimes 2}$ defined by $\Delta_{\#}(v) = 0$ on V , by $\Delta_{\#}(X) := \Delta(X)$ on X and extended by induction on $As(V)\langle X \rangle$ by,*

$$\Delta_{\#}(a_1 a_2) := \Delta_{\#}(a_1) a_2 + a_1 \Delta_{\#}(a_2), \quad \forall a_1, a_2 \in As(V)\langle X \rangle,$$

is coassociative.

Proof: Straightforward. □

Remark: [Interpretation] As a co-operation “breaks down” an element say x into pieces $x_{(1)}$ and $x_{(2)}$ in several manners, we can say that elements of V are indecomposable but not those of X .

Let G be a weighted directed graph and G_n be the set of path of length n , in particular G_0 is the vertex set and G_1 the arrow set. A directed arrow $e_i \rightarrow e_j$, with $e_i, e_j \in G_0$, will be denoted by $a_{(i,j)}$. A directed graph is said to be *weighted* if it is equipped with a weight map, i.e., with a map w from G_1 to k . The relations $e_i e_j = 1$ if $i = j$ and 0 otherwise, $a_{(i,j)} a_{(k,l)} = 0$ if $j \neq k$, $e_i a_{(k,l)} = a_{(k,l)}$ if $i = k$ and 0 otherwise and $a_{(k,l)} e_i = a_{(k,l)}$ if $i = l$ and 0 otherwise, for every $e_i, e_j \in G_0$ and $a_{(i,j)}, a_{(k,l)} \in G_1$, turn the space $kG := \bigoplus_{n=0}^{\infty} kG_n$ into a k -algebra called the path algebra. The product is then the concatenation of path whenever possible. We improve a construction given in [1] in the case of directed graphs (trivially weighted). Define the following co-operation Δ , for any $e_i \in G_0$ by $\Delta(e_i) := 0$, for any weighted arrow $a_{(i,j)}$ by $\Delta a_{(i,j)} := w(a_{(i,j)}) e_i \otimes e_j$ and for any weighted path $\alpha := a_{(i_1, i_2)} a_{(i_2, i_3)} \dots a_{(i_{n-1}, i_n)}$ by,

$$\Delta(\alpha) := \Delta(a_{(i_1, i_2)}) a_{(i_2, i_3)} \dots a_{(i_{n-1}, i_n)} + \dots + a_{(i_1, i_2)} a_{(i_2, i_3)} \dots a_{(i_{n-2}, i_{n-1})} \Delta a_{(i_{n-1}, i_n)}.$$


Proposition 2.32 ([9]) *Equipped with this co-operation, any weighted directed graph carries a ϵ -bialgebra structure.*

Remark: Elements which are unbreakable with regards to the co-operation are the vertices of the graph.

Remark: [Insertion of sequences and Virus attacks] From ϵ -bialgebras, pre-Lie algebras can be constructed. Let (A, μ, Δ) be a ϵ -bialgebra. Add a pre-Lie structure on A via the following operation [1]:

$$\alpha \bowtie \alpha' := \alpha'_{(1)} \alpha \alpha'_{(2)}, \quad \forall \alpha, \alpha' \in A, \quad \text{with } \Delta(\alpha) := \alpha_{(1)} \otimes \alpha_{(2)}.$$

Our structure on weighted directed graphs can be interpreted as follows. When virus attack a cell, it breaks its D.N.A. chain α into two pieces $\alpha_{(1)}$ and $\alpha_{(2)}$ to insert its own D.N.A. chain. The result gives the D.N.A. chain $\alpha_{(1)} \alpha \alpha_{(2)}$. However, to do that, we can say that virus need an energy λ_{12} , or breaking the D.N.A. chain α of the host cell at this place can be done with a probability λ_{12} .



Insertion of a viral D.N.A. chain into the D.N.A. of a Host cell. When all possibilities are taken into account, this insertion mechanism has a pre-Lie structure coming from a natural ϵ -bialgebra structure.

With our modification of the coproduct introduced by Aguiar on graphs, such attacks and the probability or the energy to break the D.N.A. chain at this place rather than another place can be modelled. The coproduct, viewed as a map which “reads sequentially” the D.N.A. chain, models all the possibilities to open the D.N.A. chain of the host cell with a given probability or a given energy. This entails that the insertion of a viral D.N.A. chain into the genetic code of the host cell has a natural pre-Lie structure.

This model can be improved by introducing more parameters to open a chain. This leads to the construction of n coproducts Δ_k , $k = 1, \dots, n$, such as the one defined in this example, but with the family of weights $(\lambda_{ij})_{(i \rightarrow j \in G_1)}$ replaced by other families $(\lambda_{ij}^{(k)})_{(i \rightarrow j \in G_1)}$, where $k = 1, \dots, n$. Observe then that these coproducts obey the following n^2 relations:

$$(\Delta_k \otimes id)\Delta_l := (id \otimes \Delta_l)\Delta_k,$$

for all $k, l = 1, \dots, n$. In [9], such structures have been used to produce formal deformation à la Gerstenhaber of quadri-algebras. See also Section 4. Such structures appear in [9, 12] and in free probability [22].

2.7 Other constructions

Consider the k -algebra $M_2(k)$ and the following co-operations.

$$\Delta_0\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \begin{pmatrix} 0 & a \\ 0 & c \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} c & d \\ 0 & 0 \end{pmatrix},$$

$$\Delta_1\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) = \begin{pmatrix} b & 0 \\ d & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ a & b \end{pmatrix},$$

Observe that $(M_2(k), \Delta_0)$, found in [1], and $(M_2(k), \Delta_1)$ are both ϵ -bialgebras. The same can be obtained on $M_n(k)$ by defining n coproducts $\Delta_{(a,b)}$ with $a + b = n + 1$ such that,

$$\Delta_{(a,b)}X := \sum_k X_{ka}E_{kb} \otimes E_{ab} - E_{ab} \otimes \sum_{k'} X_{bk'}E_{ak'},$$

where $X := X_{ij}E_{ij}$ with $X_{ij} \in k$ and $E_{ij} \in M_n(k)$ are the matrices with a 1 at the intersection of the row i and the column j . For every (a, b) with $a + b = n + 1$, $(M_n(k), \Delta_{(a,b)})$ is a ϵ -bialgebra.

3 Algebraic structures related to two entangled coproducts.

We continue our exploration of these particular bialgebras by considering this time k -vector spaces equipped with two co-operations.

3.1 L -Coalgebras $(A, \mu, \Delta_1, \Delta_2)$, with (A, μ, Δ_2) a $\epsilon'[R]$ -bialgebra.

Let (A, μ) be an associative algebra equipped with two different co-operations $\Delta_1, \Delta_2 : A \rightarrow A^{\otimes 2}$. The k -algebra $\text{End}(A)$ has two convolution products denoted by $*_1$ and $*_2$. In the sequel, we will use four linear maps $\beta_1, \beta_2, \gamma_1, \gamma_2 : \text{End}(A) \rightarrow \text{End}(A)$ defined by $\beta_1(T) := id *_1 T$, $\beta_2(T) := id *_2 T$, $\gamma_1(T) := T *_1 id$, $\gamma_2(T) := T *_2 id$, for all $T \in \text{End}(A)$.

Proposition 3.1 *Let $(A, \mu, \Delta_1, \Delta_2)$ be an associative algebra equipped with two co-operations Δ_1, Δ_2 . Suppose (A, μ, Δ_2) is a $\epsilon'[R]$ -bialgebra. Then,*

$$(Bax.12) \quad \beta_2(T)\beta_1(S) = \beta_1(\beta_2(T)S), \quad \forall T, S \in \text{End}(A).$$

In addition, the relations between operations labelled by 1 and 2 are stated in the following 2 by 2 matrix $(M_{4,\beta})_{ij}$. For all $T, R, S \in \text{End}(A)$,

$$\begin{aligned} R\overleftarrow{\star}_{\beta_1}(S \prec_{\beta_2} T) &= (R\overleftarrow{\star}_{\beta_1} S) \prec_{\beta_2} T, \quad R\overleftarrow{\star}_{\beta_2}(S \prec_{\beta_1} T) = (R\overleftarrow{\star}_{\beta_2} S) \prec_{\beta_1} T, \\ R \prec_{\beta_1} (S\overleftarrow{\star}_{\beta_2} T) &= (R \prec_{\beta_2} S) \prec_{\beta_1} T, \quad R\overleftarrow{\star}_{\beta_2}(S\overleftarrow{\star}_{\beta_1} T) = (R\overleftarrow{\star}_{\beta_2} S)\overleftarrow{\star}_{\beta_1} T. \end{aligned}$$

If (A, μ, Δ_1) is also a $\epsilon'[R]$ -bialgebra then $(\text{End}(A), \overleftarrow{\star}_{\beta_2} \dashrightarrow \overleftarrow{\star}_{\beta_1})$ is an associative L -algebra.

Proof: Let $(A, \mu, \Delta_1, \Delta_2)$ be an algebra equipped with two co-operations Δ_1, Δ_2 . Suppose (A, μ, Δ_2) is a $\epsilon'[R]$ -bialgebra. Fix $T, S \in \text{End}(A)$ and $x \in A$. Set $\Delta_1(x) := x_{(1)} \otimes x_{(2)}$ and $\Delta_2(x) := x^{(1)} \otimes x^{(2)}$. On the one hand,

$$\begin{aligned} \beta_2(T)\beta_1(S)(x) &:= \beta_2(T)(x_{(1)}S(x_{(2)})), \\ &= \mu(id \otimes T)\Delta_2(x_{(1)}S(x_{(2)})), \\ &= \mu(id \otimes T)(x_{(1)}S(x_{(2)})^{(1)} \otimes S(x_{(2)})^{(2)}), \\ &= x_{(1)}S(x_{(2)})^{(1)}T(S(x_{(2)})^{(2)}). \end{aligned}$$

On the other hand,

$$\begin{aligned} \beta_1(\beta_2(T)S)(x) &:= \mu(id \otimes \beta_2(T)S)\Delta_1(x), \\ &= x_{(1)}\beta_2(T)S(x_{(2)}), \\ &= x_{(1)}\mu(id \otimes T)\Delta_2(S(x_{(2)})), \\ &= x_{(1)}\mu(id \otimes T)S(x_{(2)})^{(1)} \otimes S(x_{(2)})^{(2)}, \\ &= x_{(1)}S(x_{(2)})^{(1)}T(S(x_{(2)})^{(2)}). \end{aligned}$$

Therefore,

$$(Bax.12) \quad \beta_2(T)\beta_1(S) = \beta_1(\beta_2(T)S), \quad \forall T, S \in \text{End}(A).$$

The other equalities are straightforward consequences of (Bax.12). If (A, μ, Δ_1) is also a $\epsilon'[R]$ -bialgebra then $(\text{End}(A), \overleftarrow{\star}_{\beta_1})$ is an associative algebra, see for instance Theorem 2.2. As $\overleftarrow{\star}_{\beta_2}$ is also associative and is linked to $\overleftarrow{\star}_{\beta_1}$ via the equality $(M_{4,\beta})_{22}$, the k -vector space $(\text{End}(A), \overleftarrow{\star}_{\beta_2} \dashrightarrow \overleftarrow{\star}_{\beta_1})$ is an associative L -algebra. \square

Proposition 3.2 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra, i.e., whose coproducts verify:*

$$(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1.$$

Suppose (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra. Then,

$$(Bax. 3) \quad \gamma_2(T)\gamma_1(S) = \gamma_2(T\gamma_1(S)).$$

In addition, the operations labelled by 1 and 2 are stated in the following 2 by 2 matrix $(M_{4,\gamma})_{ij}$. For all $R, S, T \in \text{End}(A)$,

$$\begin{aligned} R \succ_{\gamma_2} (S \overrightarrow{\star}_{\gamma_1} T) &= (R \succ_{\gamma_2} S) \overrightarrow{\star}_{\gamma_1} T, & R \succ_{\gamma_1} (S \overrightarrow{\star}_{\gamma_2} T) &= (R \succ_{\gamma_1} S) \overrightarrow{\star}_{\gamma_2} T, \\ R \overrightarrow{\star}_{\gamma_2} (S \overrightarrow{\star}_{\gamma_1} T) &= (R \overrightarrow{\star}_{\gamma_2} S) \overrightarrow{\star}_{\gamma_1} T, & R \succ_{\gamma_2} (S \prec_{\gamma_1} T) &= (R \overrightarrow{\star}_{\gamma_1} S) \prec_{\gamma_2} T. \end{aligned}$$

Moreover, for all $i, j = \{1, 2\}$ and $R, S, T \in \text{End}(A)$,

$$R \overleftarrow{\star}_{\beta_i} (S \overrightarrow{\star}_{\gamma_j} T) = (R \overleftarrow{\star}_{\beta_i} S) \overrightarrow{\star}_{\gamma_j} T$$

$$R \succ_{\gamma_i} (S \prec_{\beta_j} T) = (R \succ_{\gamma_i} S) \prec_{\beta_j} T.$$

If (A, μ, Δ_1) is a $\epsilon[R]$ -bialgebra then $(\text{End}(A), \overrightarrow{\star}_{\gamma_2} \dashrightarrow \overrightarrow{\star}_{\gamma_1})$ is an associative L -algebra.

Proof: Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra. Suppose (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra. Fix $T, S \in \text{End}(A)$ and $x \in A$. Set $\Delta_1(x) := x_{(1)} \otimes x_{(2)}$ and $\Delta_2(x) := x^{(1)} \otimes x^{(2)}$. On the one hand,

$$\begin{aligned} \gamma_2(T)\gamma_1(S)(x) &= \mu(T \otimes id)\Delta_2(S(x_{(1)})x_{(2)}), \\ &= \mu(T \otimes id)(S(x_{(1)})(x_{(2)})^{(1)} \otimes (x_{(2)})^{(2)}), \\ &= T(S(x_{(1)})(x_{(2)})^{(1)})(x_{(2)})^{(2)}. \end{aligned}$$

On the other hand,

$$\begin{aligned} \gamma_2(T\gamma_1(S))(x) &= \mu(T\gamma_1(S) \otimes id)\Delta_2(x), \\ &= T\gamma_1(S)(x^{(1)})x^{(2)}, \\ &= T\mu(S \otimes id)(x^{(1)})_{(1)} \otimes (x^{(1)})_{(2)}x^{(2)}, \\ &= T(S((x^{(1)})_{(1)})(x^{(1)})_{(2)})x^{(2)}, \end{aligned}$$

whence the equality Bax. 3 since $\Delta_1 \dashrightarrow \Delta_2$. The matrix $M_{4,\gamma}$ is obtained by using Bax. 3. If (A, μ, Δ_1) is a $\epsilon[R]$ -bialgebra then, $(\text{End}(A), \overrightarrow{\star}_{\gamma_1})$ is an associative algebra. As $\overrightarrow{\star}_{\gamma_1}$ and $\overrightarrow{\star}_{\gamma_2}$ are related by the equation $(M_{4,\gamma})_{21}$, the k -vector space $(\text{End}(A), \overrightarrow{\star}_{\gamma_2} \dashrightarrow \overrightarrow{\star}_{\gamma_1})$ is an associative L -algebra. \square

Proposition 3.3 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra. Suppose (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra and (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra. Then, there exists a $([M]_{ij})_{(i,j)}$ -algebra structure on $\text{End}(A)$.*

Proof: As (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra, the linear map β_1 is a right Baxter operator. As (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra the linear map γ_2 is a left Baxter operator. From $(id \otimes \Delta_2)\Delta_1 = (\Delta_1 \otimes id)\Delta_2$, we get the commutation relation $\beta_1\gamma_2 = \gamma_2\beta_1$. Apply Theorem 2.10 to conclude. \square

In terms of category, the following functor is obtained,

$$[\epsilon'[R] \dashrightarrow \epsilon[R]] - \text{Bialg.} \longrightarrow ([M]_{(ij)}),$$

where $[\epsilon'[R] \dashrightarrow \epsilon[R]] - \text{Bialg.}$ denotes the category of those L -coalgebras verifying hypotheses of Proposition 3.3.

Example 3.4 We produce an example of k -vector space $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ where both (A, μ, Δ_1) and (A, μ, Δ_2) are $\epsilon[R]$ -bialgebras. Consider the Example 2.17. Let I_2 be the identity matrix and X be any non-null 2×2 -matrix. Then, the co-operation given by $\Delta_2(M) := M \otimes X$ is coassociative and obey the identity:

$$(\Delta_1 \otimes id)\Delta_2 := (id \otimes \Delta_2)\Delta_1.$$

The k -vector space $(M_2(k), \Delta_1 \dashrightarrow \Delta_2)$ is such that both $(M_2(k), \Delta_1)$ and $(M_2(k), \Delta_2)$ are $\epsilon[R]$ -bialgebras.

3.2 Dendriform dialgebras from L -coalgebras

Let us construct dendriform algebras from non-coassociative co-operations.

Proposition 3.5 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra, i.e., such that,*

$$(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1.$$

Suppose (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra and (A, μ, Δ_2) is a ϵ -bialgebra. Define the following binary operations $\prec, \succ: \text{End}(A)^{\otimes 2} \rightarrow \text{End}(A)$ by,

$$T \prec S := \beta_1(T)\gamma_2(S) \quad \text{and} \quad T \succ S := \gamma_2\beta_1(T)S, \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \prec, \succ)$ is a dendriform dialgebra.

Proof: As (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra, the linear map β_1 is a right Baxter operator and $(\text{End}(A), \overleftarrow{*}_{\beta_1})$ is an associative algebra. As (A, μ, Δ_2) is a ϵ -bialgebra, γ_2 is a Baxter operator which commutes with β_1 since $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$. This entails that γ_2 is a Baxter operator on the associative algebra $(\text{End}(A), \overleftarrow{*}_{\beta_1})$. Therefore, $(\text{End}(A), \prec, \succ)$ is a dendriform dialgebra. \square

In terms of category, the following functor is obtained,

$$[\epsilon'[R] \dashrightarrow \epsilon] - \text{Bialg.} \longrightarrow \text{DiDend.},$$

where $[\epsilon'[R] \dashrightarrow \epsilon] - \text{Bialg.}$ denotes the category of those L -coalgebras verifying hypotheses of Proposition 3.5.

Proposition 3.6 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra i.e., such that,*

$$(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1.$$

Suppose (A, μ, Δ_1) is a ϵ -bialgebra and (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra. Define the following binary operations $\succ, \prec: \text{End}(A)^{\otimes 2} \rightarrow \text{End}(A)$ by,

$$T \succ S := \beta_1(T)\gamma_2(S) \quad \text{and} \quad T \prec S := T\beta_1(\gamma_2(S)), \quad \forall T, S \in \text{End}(A).$$

Then, $(\text{End}(A), \prec, \succ)$ is a dendriform dialgebra.

Proof: As (A, μ, Δ_1) is a ϵ -bialgebra, the linear map β_1 is a Baxter operator on $\text{End}(A)$. As (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra, the linear map γ_2 is a left Baxter operator which commutes with β_1 since $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$. This entails that β_1 is a Baxter operator on the associative algebra $(\text{End}(A), \overrightarrow{*}_{\gamma_2})$. Therefore, $(\text{End}(A), \prec, \succ)$ is a dendriform dialgebra. \square

In terms of category, the following functor is obtained,

$$[\epsilon \dashrightarrow \epsilon[R]] - \text{Bialg.} \longrightarrow \text{DiDend.},$$

where $[\epsilon \dashrightarrow \epsilon[R]] - \text{Bialg.}$ denotes the category of those L -coalgebras verifying hypotheses of Proposition 3.6.


Example 3.7 [Dendriform dialgebras from weighted directed graphs and research algorithms] Keep notation of Proposition 2.32. Recall the co-operation Δ , for any $e_i \in G_0$ is defined by $\Delta(e_i) := 0$, for any weighted arrow $a_{(i,j)}$ by $\Delta a_{(i,j)} := w(a_{(i,j)})e_i \otimes e_j$ and for any weighted path $\alpha := a_{(i_1,i_2)}a_{(i_2,i_3)} \dots a_{(i_{n-1},i_n)}$ by,

$$\Delta(\alpha) := \Delta(a_{(i_1,i_2)})a_{(i_2,i_3)} \dots a_{(i_{n-1},i_n)} + \dots + a_{(i_1,i_2)}a_{(i_2,i_3)} \dots a_{(i_{n-2},i_{n-1})}\Delta a_{(i_{n-1},i_n)}.$$

Equipped with this co-operation, any weighted directed graph carries a ϵ -bialgebra structure. For any path α define the coproduct Δ_M by $\Delta_M(\alpha) := \alpha \otimes \sum_{ij} a_{ij}$ for any path α . Equipped with this co-operation, any weighted directed graph carries a $\epsilon[R]$ -bialgebra structure. Observe that,

$$(\Delta \otimes id)\Delta_M = (id \otimes \Delta_M)\Delta.$$

The map $(\mu \otimes id)(id \otimes \Delta)\Delta_M$ can be interpreted as a research map throughout the graph G since for any path α , $(\mu \otimes id)(id \otimes \Delta)\Delta_M(\alpha) := \sum_j \lambda_{t(\alpha)j} \alpha \otimes e_j$, with $\lambda_{t(\alpha)j}$ a non-null scalar if there is an arrow $t(\alpha) \rightarrow j$ with weight $\lambda_{t(\alpha)j}$ belonging to G_1 . Consider the following weighted directed graph G , with weights λ_{ij} different from zero if $i \rightarrow j \in G_1$ and zero otherwise.



The weighted directed graph G (no weight represented).

The operator $(\mu \otimes id)(id \otimes \Delta)\Delta_M$ applied to $\alpha := a_{12}a_{23}a_{34}$ gives,

$$a_{12}a_{23}a_{34} \xrightarrow{\Delta_M} \sum_{i,j=1}^6 a_{12}a_{23}a_{34} \otimes a_{ij} \xrightarrow{id \otimes \Delta} \sum_{i,j=1}^6 \lambda_{ij} a_{12}a_{23}a_{34} \otimes e_i \otimes e_j \xrightarrow{\mu \otimes id} \lambda_{45} a_{12}a_{23}a_{34} \otimes e_5.$$

3.3 Entanglement of left and right bialgebras

Proposition 3.8 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra such that (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra and (A, μ, Δ_2) is a $\epsilon'[L]$ -bialgebra. Then, there exists a $([M_1]_{ij})_{(i,j)}$ -algebra structure on $End(A)$.*

Proof: As (A, μ, Δ_2) is a $\epsilon'[L]$ -bialgebra, γ_2 is a right Baxter operator. So is β_1 since (A, μ, Δ_1) is a $\epsilon'[R]$ -bialgebra. The two right Baxter operators commute, i.e., $\gamma_2\beta_1 = \beta_1\gamma_2$, since the notation $\Delta_1 \dashrightarrow \Delta_2$ means $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$. Consequently, γ_2 is a right Baxter operator on the L -anti-dipterous $(End(A), \overleftarrow{*}_{\beta_1}, \prec_{\beta_1})$. Corollary 2.13 claims that $End(A)$ has a structure of $([M_1]_{ij})_{(i,j)}$ -algebra. \square

In terms of category, the following functor is obtained,

$$[\epsilon'[R] \dashrightarrow \epsilon'[L]] - \mathbf{Bialg.} \longrightarrow [M_1]_{(ij)},$$

where $[\epsilon'[R] \dashrightarrow \epsilon'[L]] - \mathbf{Bialg.}$ denotes the category of those L -coalgebras verifying hypotheses of Proposition 3.8.

Proposition 3.9 *Let $(A, \mu, \Delta_1 \dashrightarrow \Delta_2)$ be a L -coalgebra such that (A, μ, Δ_1) is a $\epsilon[L]$ -bialgebra and (A, μ, Δ_2) a $\epsilon[R]$ -bialgebra. Then, there exists a $([M_2]_{ij})_{(i,j)}$ -algebra structure on $End(A)$.*

Proof: As (A, μ, Δ_2) is a $\epsilon[R]$ -bialgebra, γ_2 is a left Baxter operator. So is β_1 since (A, μ, Δ_1) is a $\epsilon[L]$ -bialgebra. The two left Baxter operators commute, i.e., $\gamma_2\beta_1 = \beta_1\gamma_2$, since the notation $\Delta_1 \dashrightarrow \Delta_2$ means $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$. Consequently, γ_2 is a left Baxter operator on the L -dipterous $(End(A), \overrightarrow{*}_{\beta_1}, \succ_{\beta_1})$. Proposition 2.26 claims that $End(A)$ has a structure of $([M_2]_{ij})_{(i,j=1,\dots,3)}$ -algebra. \square

In terms of category, the following functor is obtained,

$$[\epsilon[L] \dashrightarrow \epsilon[R]] - \mathbf{Bialg.} \longrightarrow [\mathbf{M}_2]_{(ij)},$$

where $[\epsilon[R] \dashrightarrow \epsilon[L]] - \mathbf{Bialg.}$ denotes the category of those L -coalgebras verifying hypotheses of Proposition 3.9.

4 Splitting associative clusters into associative operations

Let (X, \bullet) be a k -algebra and $(\bullet_i)_{1 \leq i \leq N} : X^{\otimes 2} \rightarrow X$ be a family of binary operations on X . Recall that $\bullet \longrightarrow \sum_i \bullet_i$ means $x \bullet y = \sum_i x \bullet_i y$, for all $x, y \in X$, i.e., \bullet *splits* into the N operations $\bullet_1, \dots, \bullet_N$ or is a *cluster of N (binary) operations*. Such a cluster will be called associative if \bullet is associative. For instance dendriform dialgebras have their associative operations which split into two operations $\star \longrightarrow \prec + \succ$. Here, we will be interested in associative clusters which split into n associative products.

4.1 n -Hypercubic $\epsilon[R]$ -bialgebras

Definition 4.1 [Hypercubic algebras] Fix a non-null integer n . A n -hypercubic algebra $(A, \star_1, \dots, \star_n)$ is a k -vector space equipped with n binary operations $\star_1, \dots, \star_n : A^{\otimes 2} \rightarrow A$, such that for all $x, y, z \in A$ and for all $i, j \in \{1, \dots, n\}$:

$$(Hyp.) \quad (x \star_i y) \star_j z = x \star_i (y \star_j z).$$

Similarly, a n -hypercubic coalgebra $(A, \Delta_1, \dots, \Delta_n)$ is a k -vector space equipped with n co-operations $\Delta_1, \dots, \Delta_n : A \rightarrow A^{\otimes 2}$, such that for all $i, j \in \{1, \dots, n\}$:

$$(CoHyp.) \quad (\Delta_i \otimes id) \Delta_j = (id \otimes \Delta_j) \Delta_i.$$

A n -hypercubic $\epsilon[R]$ -bialgebra $(A, \mu, \Delta_1, \dots, \Delta_n)$ is a n -hypercubic coalgebra $(A, \Delta_1, \dots, \Delta_n)$ together with an associative algebra (A, μ) such that for all $i = 1, \dots, n$, (A, μ, Δ_i) are $\epsilon[R]$ -bialgebras. Similarly for the left version. A n -hypercubic ϵ -bialgebra is a n -hypercubic coalgebra $(A, \Delta_1, \dots, \Delta_n)$ together with an associative algebra (A, μ) such that for all $i = 1, \dots, n$, (A, μ, Δ_i) are ϵ -bialgebras. Such bialgebras have been used in [9] to produce formal deformations of associative algebras such as dendriform di- and tri-algebras, quadri-algebras and ennea-algebras.

Remark: Observe that axioms $(Hyp.)$ entails the associativity of all the involved operations. Moreover these axioms are invariant by action of $M_n(k)$. More precisely, if $M \in M_n(k)$ and $X := (\star_1, \dots, \star_n)$ then ${}^t(M({}^tX)) := (\star'_1, \dots, \star'_n)$ will still yield associative products $\star'_1, \dots, \star'_n$ which will satisfy the axioms stated above. In particular, $\sum_{i=1}^n \star_i$ is still associative. If $(A, \star_1, \dots, \star_n)$ is a n -hypercubic algebra, then \star_j is a Hochschild 2-cocycle on the associative algebra (A, \star_i) , for all $i, j = 1, \dots, n$. In addition, there also exists a notion of opposite structure. Indeed, define new operations $x \star_i^{op} y := y \star_i x$, for all $x, y \in A$ and $i = 1, \dots, n$ and observe that $(A, \star_1^{op}, \dots, \star_n^{op})$ is still an hypercubic algebra.

Remark: Since the operad associated with hypercubic algebras is binary, quadratic and non-symmetric, the free n -hypercubic algebra on a k -vector space V is given by:

$$H^n(V) := \bigoplus_{p \geq 1} (H^n)_p \otimes V^{\otimes p}.$$

On one generator, one see that $H^n(k) := \bigoplus_{p \geq 1} (H^n)_p$. The dimension of $(H^n)_p$ is n^p . These operads are conjectured to be Koszul self-dual. This has already been proved in the case $p = 2$ [20] and $p = 3$ [17].

Let us construct hypercubic algebras from non-coassociative coproducts.

Theorem 4.2 *Let $(A, \mu, \Delta_1, \dots, \Delta_n)$ be a k -vector space such that for all $i = 1, \dots, n$, (A, μ, Δ_i) are $\epsilon'[R]$ -bialgebras. Then, $(\text{End}(A), \overleftarrow{*}_{\beta_1}, \dots, \overleftarrow{*}_{\beta_n})$ is a n -hypercubic algebra.*

Proof: As for all $i = 1, \dots, n$, (A, μ, Δ_i) are $\epsilon'[R]$ -bialgebras, $(\text{End}(A), \overleftarrow{*}_{\beta_i})$ are associative algebras. Apply now the last claim of Proposition 3.1 to any pair $((A, \mu, \Delta_i), (A, \mu, \Delta_j))$ to prove that $(\text{End}(A), \overleftarrow{*}_{\beta_1}, \dots, \overleftarrow{*}_{\beta_n})$ is a n -hypercubic algebra. \square

Proposition 4.3 *Let $(A, \mu, \Delta_1, \dots, \Delta_n)$ be a n -hypercubic $\epsilon[R]$ -bialgebra. Then, the k -vector space $(\text{End}(A), \overrightarrow{*}_{\gamma_1}, \dots, \overrightarrow{*}_{\gamma_n})$ is a n -hypercubic algebra.*

Proof: Apply now the last claim of Proposition 3.2 to any pair $((A, \mu, \Delta_i), (A, \mu, \Delta_j))$. \square

Proposition 4.4 *Let $(A, \mu, \Delta_1, \Delta_2)$ be a 2-hypercubic ϵ -bialgebra. Then, the associative products $\overleftarrow{*}_{\beta_1}$ and $\overleftarrow{*}_{\beta_2}$ defined on $\text{End}(A)$ verify the relation:*

$$(Aas.): \quad (R *_{\beta_1} S) *_{\beta_2} T + (R *_{\beta_2} S) *_{\beta_1} T = R *_{\beta_1} (S *_{\beta_2} T) + R *_{\beta_2} (S *_{\beta_1} T).$$

*Moreover the operations $\star_{[\lambda_1, \lambda_2]} := \lambda_1(*_{\beta_1}) + \lambda_2(*_{\beta_2}) : \text{End}(A) \rightarrow \text{End}(A)$, where $\lambda_1, \lambda_2 \in k$ are still associative.*

Proof: Let $(A, \mu, \Delta_1, \Delta_2)$ be a 2-hypercubic ϵ -bialgebra. Fix $R, S, T \in \text{End}(A)$. Observe that,

$$\beta_1(T)\beta_2(S) := \beta_1(T\beta_2(S)) + \beta_2(\beta_1(T)S),$$

if $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$. This gives (Aas') :

$$(R *_{\beta_1} S) *_{\beta_2} T - R *_{\beta_1} (S *_{\beta_2} T) = \beta_2(R\beta_1(S))T + R\beta_2(\beta_1(S)T) - R\beta_1(\beta_2(S)T) - \beta_1(R\beta_2(S))T.$$

The equation $(Aas.)$ holds by noticing that (Aas') is anti-symmetric with regards to the permutation $(2, 1)$. The last assertion is straightforward. \square

Remark: The condition $(Aas.)$ claims that $*_{\beta_1}$ is a Hochschild 2-cocycle on the associative algebra $(\text{End}(A), *_{\beta_2})$ or that $*_{\beta_2}$ is a Hochschild 2-cocycle on the associative algebra $(\text{End}(A), *_{\beta_1})$.

Remark: Observe here that associative products $\star_{[\lambda_1, \lambda_2]} \rightarrow \lambda_1(\overleftarrow{*}_{\beta_1}) + \lambda_2(\overleftarrow{*}_{\beta_2})$ split into two associative ones. The following corollary improves this result.

Corollary 4.5 *Let $(A, \mu, \Delta_1, \dots, \Delta_n)$ be a n -hypercubic ϵ -bialgebra. Then, the associative products $\star := \sum_i \lambda_i \overleftarrow{\star}_{\beta_i}$, $\hat{\star} := \sum_i \lambda_i' \overrightarrow{\star}_{\gamma_i} : \text{End}(A) \rightarrow \text{End}(A)$ are associative.*

Proof: Straightforward. For instance, fix $R, S, T \in \text{End}(A)$. Observe that,

$$\gamma_i(T)\gamma_j(S) := \gamma_i(T\gamma_j(S)) + \gamma_j(\gamma_i(T)S),$$

if $(\Delta_j \otimes \text{id})\Delta_i = (\text{id} \otimes \Delta_i)\Delta_j$. With β replaced by γ , the relation (Aas.) still holds. Therefore $\hat{\star} := \sum_i \lambda_i' \overrightarrow{\star}_{\gamma_i}$ is associative. \square

Remark: Solving equation (Aas.) entails the study of at least two types of algebras, the n -hypercubic algebras we have just seen and the n -circular algebras. Let us see the interest of such algebras and their co-versions.

4.2 Circular algebras

Definition 4.6 [circular algebras] A circular algebra $(A, \star_1, \dots, \star_n)$ is a k -vector space equipped with n binary operations $\star_1, \dots, \star_n : A^{\otimes 2} \rightarrow A$, such that for all $x, y, z \in A$ and for all $i, j \in \{1, \dots, n\}$:

$$(x \star_i y) \star_j z = x \star_j (y \star_i z).$$

Remark: Observe that such a definition entails the associativity of all the involved operations and that axioms of a circular algebra are invariant by action of $M_n(k)$. More precisely, if $M \in M_n(k)$ and $X := (\star_1, \dots, \star_n)$ then ${}^t(M({}^tX)) := (\star'_1, \dots, \star'_n)$ will still yield associative products $\star'_1, \dots, \star'_n$ which will satisfy the axioms stated above. In particular, $\sum_{i=1}^n \star_i$ is still associative.

Remark: If $(A, \star_1, \dots, \star_n)$ is a n -circular algebra, then \star_j is a Hochschild 2-cocycle on the associative algebra (A, \star_i) .

Remark: If there exists a common unit to all these products \star_1, \dots, \star_n , then they are equal pairwise.

Remark: Since the non-symmetric operad associated with circular algebras is binary and quadratic, the free circular algebra on a k -vector space V is given by:

$$C^n(V) := \bigoplus_{p \geq 1} (C^n)_p \otimes V^{\otimes p}.$$

On one generator, one see that $C^n(k) := \bigoplus_{p \geq 1} (C^n)_p$. The dimension of C_p is n^p .

4.2.1 Constructions of n -circular algebras via right Baxter operators

Lemma 4.7 *Let (A, μ) be an associative algebra. Suppose there exist n linear maps $\phi_1, \dots, \phi_n : A \rightarrow A$ such that for all $x, y \in A$ and for $i, j \in \{1, \dots, n\}$,*

$$\phi_i(x)\phi_j(y) := \phi_i(\phi_j(xy)),$$

hold. Define the binary operations \star_1, \dots, \star_n by $x \star_1 y := \phi_1(x)y, \dots, x \star_n y := \phi_n(x)y$. Then, the k -vector space $(A, \star_1, \dots, \star_n)$ is a n -circular algebra. Similarly, Suppose these n -linear maps verify,

$$\phi_i(x)\phi_j(y) := \phi_j(x\phi_i(y)),$$

for $i, j \in \{1, \dots, n\}$. Then, the binary operations $\star'_1, \dots, \star'_n$ define by $x \star'_1 y := x\phi_1(y), \dots, x \star'_n y := x\phi_n(y)$ turn the k -vector space $(A, \star'_1, \dots, \star'_n)$ into a n -circular algebra.

Proof: Straightforward. □

Proposition 4.8 *Let $(A, \mu, \Delta_1, \dots, \Delta_n)$ be a n -circular $\epsilon[R]$ -bialgebra. Then, there exists a n -circular algebraic structure on $\text{End}(A)$.*

Proof: Recall that $\gamma_i(T) := T *_{\epsilon_i} id$, for all $T \in \text{End}(A)$ and $i = 1, \dots, n$. Fix $T, S \in \text{End}(A)$ and $a \in A$. Set $\Delta_1(a) := a_{(1)} \otimes a_{(2)}$ and $\Delta_1(a) := a^{(1)} \otimes a^{(2)}$. On the one hand, $\gamma_2(T)\gamma_1(S)(a) := T(S(a_{(1)})(a_{(2)})^{(1)})(a_{(2)})^{(2)}$. On the other hand, $\gamma_1(T\gamma_2(S))(a) := T\mu(S \otimes id)\Delta_2(a_{(1)})a_{(2)} = T(S((a_{(1)})^{(1)})(a_{(1)})^{(2)})a_{(2)}$. The two relations are equal if $\Delta_2(a_{(1)}) \otimes a_{(2)} = a_{(1)} \otimes \Delta_2(a_{(2)})$. We obtain a family of operators $\gamma_1, \dots, \gamma_n$, verifying for all $i, j = 1, \dots, n$, and for all $T, S \in \text{End}(A)$,

$$\gamma_i(T)\gamma_j(S) = \gamma_j(T\gamma_i(S)).$$

Apply now Lemma 4.7 to conclude. □

4.2.2 Constructions of circular algebras

We give two examples of 2-circular algebras constructed from convolution products on the k -algebra $M_2(k)$. Fix $\lambda, \nu \in k$ and consider the following co-operations.

$$\begin{aligned} \Delta_1\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) &= \begin{pmatrix} 0 & a \\ 0 & c \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} c & d \\ 0 & 0 \end{pmatrix}, \\ \Delta_2\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) &= \begin{pmatrix} 0 & \lambda a \\ 0 & \nu c \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \otimes \begin{pmatrix} \nu c & \lambda d \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

Then, $(\Delta_i \otimes id)\Delta_j = (id \otimes \Delta_i)\Delta_j$ holds for all $i, j := 1, 2$. Similarly,

$$\begin{aligned} \Delta'_1\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) &= \begin{pmatrix} b & 0 \\ d & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ a & b \end{pmatrix}, \\ \Delta'_2\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right) &= \begin{pmatrix} \lambda b & 0 \\ \nu d & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} - \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \otimes \begin{pmatrix} 0 & 0 \\ \nu a & \lambda b \end{pmatrix}. \end{aligned}$$

The equalities $(\Delta'_i \otimes id)\Delta'_j = (id \otimes \Delta'_i)\Delta'_j$ hold for all $i, j := 1, 2$. The k -algebra $M_2(k)$ equipped with the two convolution products $*_1$ and $*_2$ or \star'_1 and \star'_2 is a 2-circular algebra. We do not know how to construct 2-circular ϵ -bialgebras or their left and right versions. However, observe that $(M_2(k), \Delta_1)$ is a ϵ -bialgebra [1]. See also Subsection 2.7.

5 Octo-algebras

Up to now, we have handled two commuting Baxter operators. What kind of structures emerge if we work with three pairwise commuting Baxter operators?

5.1 Three different ways to obtain quadri-algebras from an associative cluster

We weak hypotheses of Section 4 and introduce the L -circularity. Let V be a k -vector space equipped with two co-operations Δ_1 and Δ_2 . The co-operation Δ_1 is said to be L -circular with regards to Δ_2 if:

1. $\Delta_1 \dashrightarrow \Delta_2$,
2. $(\Delta_1 \otimes id)\Delta_2 := (id \otimes \Delta_1)\Delta_2$.

This condition will be denoted by the symbol $\Delta_1 \curvearrowright \Delta_2$. One of the interest of L -circularity is to produce three Baxter operators which commute pairwise in the case of a L -circular ϵ -bialgebra, which is simply a k -vector space $(A, \mu, \Delta_1 \curvearrowright \Delta_2)$ where both (A, μ, Δ_1) and (A, μ, Δ_2) are ϵ -bialgebras. This gives birth to octo-algebras, which are simply associative algebras whose associative products are clusters of 8 operations related by 27 conditions. From such an octo-algebra, 3 quadri-algebras or 6 dendriform dialgebras sharing the same associative operation are constructed. Let us see how it works.

Definition 5.1 [Octo-algebra] An *octo-algebra* O is a k -vector space equipped with 8 binary operations:

$$\swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2, \searrow_1, \searrow_2, \nearrow_1, \nearrow_2: O^{\otimes 2} \rightarrow O.$$

To ease notation, the following sums are introduced.

$$\begin{aligned} x \prec_i y &:= x \swarrow_i y + x \nwarrow_i y; & x \bigwedge y &:= x \wedge_1 y + x \wedge_2 y; \\ x \succ_i y &:= x \searrow_i y + x \nearrow_i y; & x \bigvee y &:= x \vee_1 y + x \vee_2 y, \\ x \wedge_i y &:= x \nwarrow_i y + x \nearrow_i y; & x \ll y &:= x \prec_1 y + x \prec_2 y, \\ x \vee_i y &:= x \swarrow_i y + x \searrow_i y; & x \gg y &:= x \succ_1 y + x \succ_2 y, \\ x \circ_{12} y &:= x \circ_1 y + x \circ_2 y, & \text{with } \circ \in \{ \nearrow, \searrow, \swarrow, \nwarrow \}, \\ x \Sigma_1 y &:= x \vee_1 y + x \wedge_1 y = x \prec_1 y + x \succ_1 y; & x \Sigma_2 y &:= x \vee_2 y + x \wedge_2 y = x \prec_2 y + x \succ_2 y; \end{aligned}$$

and,

$$x \bar{\star} y := x \Sigma_1 y + x \Sigma_2 y.$$

We present our 27 relations in the following 9×3 matrix $[O_{ij}]_{(i:=1,\dots,9; j:=1,\dots,3)}$. This matrix has 9 blocs of three rows denoted by $(X_{ij})_{(i,j:=1,\dots,3)}$. For instance, X_{11} is the 3×1 -matrix $([O]_{11}, [O]_{21}, [O]_{31})$, X_{12} is the 3×1 -matrix $([O]_{12}, [O]_{22}, [O]_{32})$, and so forth.

$$\begin{aligned}
(x \nearrow_1 y) \nwarrow_1 z &= x \nwarrow_1 (y \bar{\star} z); & (x \nearrow_1 y) \nwarrow_1 z &= x \nearrow_1 (y \ll z); & (x \wedge_1 y) \nearrow_1 z &= x \nearrow_1 (y \gg z); \\
(x \swarrow_1 y) \nwarrow_1 z &= x \swarrow_1 (y \wedge z); & (x \swarrow_1 y) \nwarrow_1 z &= x \swarrow_1 (y \nwarrow_{12} z); & (x \vee_1 y) \nearrow_1 z &= x \swarrow_1 (y \nearrow_{12} z); \\
(x \prec_1 y) \swarrow_1 z &= x \swarrow_1 (y \vee z); & (x \succ_1 y) \swarrow_1 z &= x \swarrow_1 (y \swarrow_{12} z); & (x \Sigma_1 y) \swarrow_1 z &= x \swarrow_1 (y \swarrow_{12} z);
\end{aligned}$$

$$\begin{aligned}
(x \nwarrow_2 y) \nwarrow_1 z &= x \nwarrow_2 (y \Sigma_1 z); & (x \nearrow_2 y) \nwarrow_1 z &= x \nearrow_2 (y \prec_1 z); & (x \wedge_2 y) \nearrow_1 z &= x \nearrow_2 (y \succ_1 z); \\
(x \swarrow_2 y) \nwarrow_1 z &= x \swarrow_2 (y \wedge_1 z); & (x \swarrow_2 y) \nwarrow_1 z &= x \swarrow_2 (y \nwarrow_1 z); & (x \vee_2 y) \nearrow_1 z &= x \swarrow_2 (y \nearrow_1 z); \\
(x \prec_2 y) \swarrow_1 z &= x \swarrow_2 (y \vee_1 z); & (x \succ_2 y) \swarrow_1 z &= x \swarrow_2 (y \swarrow_1 z); & (x \Sigma_2 y) \swarrow_1 z &= x \swarrow_2 (y \swarrow_1 z);
\end{aligned}$$

$$\begin{aligned}
(x \nwarrow_{12} y) \nwarrow_2 z &= x \nwarrow_2 (y \Sigma_2 z); & (x \nearrow_{12} y) \nwarrow_2 z &= x \nearrow_2 (y \prec_2 z); & (x \wedge y) \nearrow_2 z &= x \nearrow_2 (y \succ_2 z); \\
(x \swarrow_{12} y) \nwarrow_2 z &= x \swarrow_2 (y \wedge_2 z); & (x \swarrow_{12} y) \nwarrow_2 z &= x \swarrow_2 (y \nwarrow_2 z); & (x \vee y) \nearrow_2 z &= x \swarrow_2 (y \nearrow_2 z); \\
(x \ll y) \swarrow_2 z &= x \swarrow_2 (y \vee_2 z); & (x \gg y) \swarrow_2 z &= x \swarrow_2 (y \swarrow_2 z); & (x \bar{\star} y) \swarrow_2 z &= x \swarrow_2 (y \swarrow_2 z).
\end{aligned}$$

Theorem 5.2 *Keep notation of Definition 5.1. If O is an octo-algebra then there exist three quadri-algebras on O ,*

1. *The depth structure, $(O_d, \nearrow_{12}, \swarrow_{12}, \swarrow_{12}, \nwarrow_{12})$,*
2. *The vertical structure, $(O_v, \prec_1, \prec_2, \succ_1, \succ_2)$,*
3. *The horizontal structure, $(O_h, \vee_1, \vee_2, \wedge_1, \wedge_2)$,*


with:

$$\begin{aligned}
x \bar{\star} y &= x \nearrow_{12} y + x \swarrow_{12} y + x \swarrow_{12} y + x \nwarrow_{12} y, \\
&= x \prec_1 y + x \prec_2 y + x \succ_1 y + x \succ_2 y, \\
&= x \vee_1 y + x \vee_2 y + x \wedge_1 y + x \wedge_2 y.
\end{aligned}$$

The rôles of the different binary operations are indicated in the following dictionary. If $(Q, \nearrow, \swarrow, \swarrow, \nwarrow, (\prec, \succ, \vee, \wedge))$ denotes a quadri-algebra obeying thus axioms of Definition 2.29, then:

1. *For the quadri-algebra $(O_d, \nearrow_{12}, \swarrow_{12}, \swarrow_{12}, \nwarrow_{12})$, we get the equivalences:*
 $\nearrow_{12} \equiv \nearrow, \swarrow_{12} \equiv \swarrow, \swarrow_{12} \equiv \swarrow, \nwarrow_{12} \equiv \nwarrow$ and $\ll \equiv \prec, \gg \equiv \succ, \vee \equiv \vee, \wedge \equiv \wedge$.
2. *For the quadri-algebra $(O_v, \prec_1, \prec_2, \succ_1, \succ_2)$, we have:*
 $\prec_1 \equiv \nearrow, \prec_2 \equiv \swarrow, \prec_2 \equiv \swarrow, \prec_1 \equiv \nwarrow$ and $\ll \equiv \prec, \gg \equiv \succ, \Sigma_2 \equiv \vee, \Sigma_1 \equiv \wedge$.
3. *For the quadri-algebra $(O_h, \vee_1, \vee_2, \wedge_1, \wedge_2)$, we obtain:*
 $\wedge_2 \equiv \nearrow, \vee_2 \equiv \swarrow, \vee_1 \equiv \swarrow, \wedge_1 \equiv \nwarrow$ and $\Sigma_1 \equiv \prec, \Sigma_2 \equiv \succ, \vee \equiv \vee, \wedge \equiv \wedge$.

Proof: Tedious, but no difficulty. Recall the 9×3 matrix $[O]$ is a 3×3 bloc matrix X with each $X_{i,j}$ denoting a bloc composed with three equations.



The depth structure is then obtained by summing as indicated in the above picture. The relations are:

$$(O_d)_{ij} := \sum_{k=1}^3 (X_{kj})_i.$$

The vertical structure is obtained by the following relations:

$$(O_v)_{ij} := \sum_{k=1}^3 (X_{ij})_k.$$

The horizontal structure is obtained by the following relations:

$$(O_h)_{ij} := \sum_{k=1}^3 (X_{ik})_j.$$

Up to a permutation of S_9 , these matrices of relations are the same that the one in Definition 2.29. \square

Remark: In terms of categories, Theorem 5.2 implies that the following diagram commutes:

$$\begin{array}{ccccc}
 \text{As.} & \xleftarrow{f} & \text{Quadri.} & \xrightarrow{f} & \text{As.} \\
 \uparrow f & \nwarrow & \uparrow F_d & \nearrow & \uparrow f \\
 \text{Quadri.} & \xleftarrow{F_h} & \text{Octo.} & \xrightarrow{F_v} & \text{Quadri.} \\
 \searrow f & \downarrow & \swarrow f & & \\
 & \text{As.} & & &
 \end{array}$$

Proposition 5.3 [Opposite of an octo-algebra] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$x \nearrow_1^{op} y := y \swarrow_2 x, \quad x \nearrow_2^{op} y := y \swarrow_1 x, \quad x \searrow_1^{op} y := y \nwarrow_2 x, \quad x \searrow_2^{op} y := y \nwarrow_1 x,$$

$$x \swarrow_1^{op} y := y \nearrow_2 x, \quad x \swarrow_2^{op} y := y \nearrow_1 x, \quad x \searrow_1^{op} y := y \searrow_2 x, \quad x \searrow_2^{op} y := y \searrow_1 x.$$

Then, for all $i = 1, 2$ we have (the labels being understood mod 2):

$$x \prec_i^{op} y := y \succ_{i+1} x, \quad x \succ_i^{op} y := y \prec_{i+1} x, \quad x \wedge_i^{op} y := y \vee_{i+1} x, \quad x \vee_i^{op} y := y \wedge_{i+1} x.$$

In addition, $O(\nearrow_1^{op}, \nearrow_2^{op}, \searrow_1^{op}, \searrow_2^{op}, \swarrow_1^{op}, \swarrow_2^{op}, \nwarrow_1^{op}, \nwarrow_2^{op})$ is an octo-algebra called the opposite of the octo-algebra $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$.

Proof: This comes from the central symmetry of center $[O]_{52} := (x \searrow_2 y) \nwarrow_1 z = x \searrow_2 (y \nwarrow_1 z)$. \square

Remark: An octo-algebra coinciding with its opposite is called a *commutative octo-algebra*. Indeed, for all x, y belonging to a commutative octo-algebra, $x \bar{x} y = y \bar{x} x$ holds.

There exist 5 other symmetries letting the matrix of relations of an octo-algebra globally invariant.

Proposition 5.4 [Transpose of an octo-algebra] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$x \nearrow_1^t y := x \swarrow_1 y, \quad x \nearrow_2^t y := x \swarrow_2 y, \quad x \swarrow_1^t y := x \nearrow_1 y, \quad x \swarrow_2^t y := x \nearrow_2 y,$$

and $x \diamond^t y := x \diamond y$ for $\diamond = \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2$. Then, for all $i = 1, 2$, we have:

$$x \prec_i^t y := x \wedge_i y, \quad x \succ_i^t y := x \vee_i y, \quad x \wedge_i^t y := x \prec_i y, \quad x \vee_i^t y := x \succ_i y,$$

and,

$$x \searrow_{12}^t y := x \searrow_{12} y, \quad x \swarrow_{12}^t y := x \nearrow_{12} y, \quad x \nearrow_{12}^t y := x \swarrow_{12} y, \quad x \nwarrow_{12}^t y := x \nwarrow_{12} y.$$

In addition, $O(\nearrow_1^t, \nearrow_2^t, \searrow_1^t, \searrow_2^t, \swarrow_1^t, \swarrow_2^t, \nwarrow_1^t, \nwarrow_2^t)$ is an octo-algebra called the transpose of the octo-algebra $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$.

Proof: Observe that $[O]_{ij} = [O^t]_{ji}$ on each 3×3 -matrices given by $(X_{11}, X_{12}, X_{13}), (X_{21}, X_{22}, X_{23})$ and (X_{31}, X_{32}, X_{33}) . \square

Proposition 5.5 [Symmetry Sym_a] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$x \searrow_1^a y := x \nearrow_2 y, \quad x \nearrow_2^a y := x \searrow_1 y, \quad x \swarrow_1^a y := x \nwarrow_2 y, \quad x \nwarrow_2^a y := x \swarrow_1 y,$$

and let the other ones fixed. Then,

$$x \searrow_{12}^a y = x \succ_2 y, \quad x \nearrow_{12}^a y = x \succ_1 y, \quad x \swarrow_{12}^a y = x \prec_2 y, \quad x \nwarrow_{12}^a y = x \prec_1 y,$$

and,

$$x \vee_2^a y := x \vee_2 y, \quad x \wedge_1^a y := x \wedge_2 y, \quad x \wedge_1^a y := x \wedge_1 y, \quad x \wedge_2^a y := x \wedge_1 y.$$

In addition, $Sym_a(O) := O(\nearrow_1^a, \nearrow_2^a, \searrow_1^a, \searrow_2^a, \swarrow_1^a, \swarrow_2^a, \nwarrow_1^a, \nwarrow_2^a)$ is an octo-algebra.

Proof: Straightforward. □

Proposition 5.6 [*Symmetry* Sym_b] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$\begin{aligned} x \searrow_1^b y &:= x \nearrow_2 y, & x \nearrow_2^b y &:= x \swarrow_2 y, & x \swarrow_1^b y &:= x \nearrow_1 y, & x \nwarrow_2^b y &:= x \swarrow_1 y, \\ x \nearrow_1^b y &:= x \nwarrow_2 y, & x \swarrow_2^b y &:= x \searrow_1 y, \end{aligned}$$

and let the two other ones fixed. Then,

$$x \searrow_{12}^b y = x \succ_2 y, \quad x \nearrow_{12}^b y = x \prec_2 y, \quad x \swarrow_{12}^b y = x \succ_1 y, \quad x \nwarrow_{12}^b y = x \prec_1 y,$$

and,

$$x \vee_2^b y := x \searrow_{12} y, \quad x \vee_1^b y := x \nearrow_{12} y, \quad x \wedge_1^b y := x \nwarrow_{12} y, \quad x \wedge_2^b y := x \swarrow_{12} y,$$

and,

$$x \prec_2^b y := x \vee_1 y, \quad x \prec_1^b y := x \wedge_1 y, \quad x \succ_1^b y := x \wedge_2 y, \quad x \succ_2^b y := x \vee_2 y.$$

In addition, $Sym_b(O) := O(\nearrow_1^b, \nearrow_2^b, \searrow_1^b, \searrow_2^b, \swarrow_1^b, \swarrow_2^b, \nwarrow_1^b, \nwarrow_2^b)$ is an octo-algebra.

Proof: Straightforward. □

Proposition 5.7 [*Symmetry* Sym_c] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$\begin{aligned} x \searrow_1^c y &:= x \swarrow_2 y, & x \nearrow_2^c y &:= x \searrow_1 y, & x \swarrow_1^c y &:= x \nwarrow_2 y, & x \nwarrow_2^c y &:= x \nearrow_1 y, \\ x \nearrow_1^c y &:= x \swarrow_1 y, & x \swarrow_2^c y &:= x \nearrow_2 y, \end{aligned}$$

and let the two other ones fixed. Then,

$$x \searrow_{12}^c y = x \vee_2 y, \quad x \nearrow_{12}^c y = x \vee_1 y, \quad x \swarrow_{12}^c y = x \wedge_2 y, \quad x \nwarrow_{12}^c y = x \wedge_1 y,$$

and,

$$x \vee_2^c y := x \succ_2 y, \quad x \vee_1^c y := x \prec_2 y, \quad x \wedge_1^c y := x \prec_1 y, \quad x \wedge_2^c y := x \succ_1 y,$$

and,

$$x \prec_2^c y := x \nearrow_{12} y, \quad x \prec_1^c y := x \nwarrow_{12} y, \quad x \succ_1^c y := x \swarrow_{12} y, \quad x \succ_2^c y := x \searrow_{12} y.$$

In addition, $Sym_c(O) := O(\nearrow_1^c, \nearrow_2^c, \searrow_1^c, \searrow_2^c, \swarrow_1^c, \swarrow_2^c, \nwarrow_1^c, \nwarrow_2^c)$ is an octo-algebra.

Proof: Straightforward. □

Proposition 5.8 [*Symmetry* Sym_d] Let $O(\nearrow_1, \nearrow_2, \searrow_1, \searrow_2, \swarrow_1, \swarrow_2, \nwarrow_1, \nwarrow_2)$ be an octo-algebra. For all $x, y \in O$, define the following binary operations as follows:

$$x \searrow_1^d y := x \swarrow_2 y, \quad x \nearrow_1^d y := x \nwarrow_2 y, \quad x \swarrow_2^d y := x \searrow_1 y, \quad x \nwarrow_2^d y := x \nearrow_1 y,$$

and let the other ones fixed. Then,

$$x \searrow_{12}^d y = x \vee_2 y, \quad x \nearrow_{12}^d y = x \wedge_2 y, \quad x \swarrow_{12}^d y = x \vee_1 y, \quad x \nwarrow_{12}^d y = x \wedge_1 y,$$

and,

$$x \vee_2^d y := x \searrow_{12} y, \quad x \vee_1^d y := x \swarrow_{12} y, \quad x \wedge_1^d y := x \nwarrow_{12} y, \quad x \wedge_2^d y := x \nearrow_{12} y,$$

and,

$$x \prec_2^d y := x \succ_1 y, \quad x \prec_1^d y := x \prec_1 y, \quad x \succ_1^d y := x \prec_2 y, \quad x \succ_2^d y := x \succ_2 y.$$

In addition, $Sym_d(O) := O(\nearrow_1^d, \nearrow_2^d, \searrow_1^d, \searrow_2^d, \swarrow_1^d, \swarrow_2^d, \nwarrow_1^d, \nwarrow_2^d)$ is an octo-algebra.

Proof: Straightforward. □

5.2 The augmented free octo-algebra and connected Hopf algebra

Let V be a k -vector space. The free octo-algebra $\mathcal{O}(V)$ on V is by definition an octo-algebra equipped with a map $i : V \mapsto \mathcal{O}(V)$ which satisfies the following universal property: for any linear map $f : V \rightarrow A$, where A is an octo-algebra, there exists a unique octo-algebra morphism $\bar{f} : \mathcal{O}(V) \rightarrow A$ such that $\bar{f} \circ i = f$.

The operad associated with octo-algebras is binary, quadratic and non-symmetric, thus of the form:

$$\mathcal{O}(V) := \bigoplus_{n \geq 1} \mathcal{O}_n \otimes V^{\otimes n}.$$

In particular, the free octo-algebra on one generator x is $\mathcal{O}(k) := \bigoplus_{n \geq 1} \mathcal{O}_n$, where $\mathcal{O}_1 := kx$, $\mathcal{O}_2 := k(x \nwarrow_1 x) \oplus k(x \nwarrow_2 x) \oplus k(x \swarrow_1 x) \oplus k(x \swarrow_2 x) \oplus k(x \nearrow_1 x) \oplus k(x \nearrow_2 x) \oplus k(x \searrow_1 x) \oplus k(x \searrow_2 x)$. The space of three variables made out of eight operations is of dimension $2 \times 8^2 = 128$. As we have 27 relations the k -vector space \mathcal{E}_3 has a dimension equal to $128 - 27 = 101$. Therefore, the sequence associated with the dimensions of $(\mathcal{E}_n)_{n \in \mathbb{N}}$ starts with 1, 8, 101... Finding the free octo-algebra on one generator is an open problem. However, a little bit more can be said on the augmented free octo-algebra.

To be as self-contained as possible, we introduce some notation to expose a theorem due to Loday [14]. Recall that a bialgebra $(H, \mu, \Delta, \eta, \kappa)$ is a unital associative algebra (H, μ, η) together with co-unital coassociative coalgebra (H, Δ, κ) . Moreover, it is required that the coproduct Δ and the counit κ are morphisms of unital algebras. A bialgebra is connected if there exists a filtration $(F_r H)_r$ such that $H = \bigcup_r F_r H$, where $F_0 H := k1_H$ and for all r ,

$$F_r H := \{x \in H; \Delta(x) - 1_H \otimes x - x \otimes 1_H \in F_{r-1} H \otimes F_{r-1} H\}.$$

Such a bialgebra admits an antipode. Consequently, connected bialgebras are connected Hopf algebras.

Let P be a binary quadratic operad. By a *unit action* [14], we mean the choice of two linear applications:

$$v : P(2) \rightarrow P(1), \quad \varpi : P(2) \rightarrow P(1),$$

giving sense, when possible, to $x \circ 1$ and $1 \circ x$, for all operations $\circ \in P(2)$ and for all x in the P -algebra A , i.e., $x \circ 1 = v(\circ)(x)$ and $1 \circ x = \varpi(\circ)(x)$. If $P(2)$ contains an associative operation, say \star , then we will require that $x \star 1 := x := 1 \star x$, i.e., $v(\star) := Id := \varpi(\star)$. We say that the *unit action* or the couple (v, ϖ) is *compatible* with the relations of the P -algebra A if they still hold on $A_+ := k1 \oplus A$. Let A, B be two P -algebras. Using the couple (v, ϖ) , we extend binary operations $\circ \in P(2)$ to the k -vector space $A \otimes 1.k \oplus k.1 \otimes B \oplus A \otimes B$ by requiring:

$$(a \otimes b) \circ (a' \otimes b') := (a \star a') \otimes (b \circ b') \quad \text{if } b \otimes b' \neq 1 \otimes 1, \quad (1)$$

$$(a \otimes 1) \circ (a' \otimes 1) := (a \circ a') \otimes 1, \quad \text{otherwise.} \quad (2)$$

The couple (v, ϖ) is said to be *coherent* with the relations of P if $A \otimes 1.k \oplus k.1 \otimes B \oplus A \otimes B$, equipped with these operations is still a P -algebra. Observe that a necessary condition for having coherence is compatibility.

One of the main interest of these two concepts is the construction of a connected Hopf algebra on the augmented free P -algebra.

Theorem 5.9 ([14]) *Let P be a binary quadratic operad. Suppose there exists an associative operation in $P(2)$. Any unit action coherent with the relations of P equips the augmented free P -algebra $P(V)_+$ on a k -vector space V with a coassociative coproduct $\Delta : P(V)_+ \rightarrow P(V)_+ \otimes P(V)_+$, which is a P -algebra morphism. Moreover, $P(V)_+$ is a connected Hopf algebra.*

Proof: See [14] for the proof. However, we reproduce it to make things clearer.

Let V be a k -vector space and $P(V)$ be the free P -algebra on V . Thanks to the coherence of the unit action, $P(V)_+ \otimes P(V)_+$ is a P -algebra. Consider the linear map $\delta : V \rightarrow P(V)_+ \otimes P(V)_+$, given by $v \mapsto 1 \otimes v + v \otimes 1$. Since, $P(V)$ is the free P -algebra on V , there exists a unique extension of δ to a morphism of augmented P -algebra $\Delta : P(V)_+ \rightarrow P(V)_+ \otimes P(V)_+$. Now, Δ is coassociative since the morphisms $(\Delta \otimes id)\Delta$ and $(id \otimes \Delta)\Delta$ extend the linear map $V \rightarrow P(V)_+^{\otimes 3}$ which maps v to $1 \otimes 1 \otimes v + 1 \otimes v \otimes 1 + v \otimes 1 \otimes 1$. By unicity of the extension, the coproduct Δ is coassociative. The bialgebra so obtained is connected. Indeed, by definition, the free P -algebra $P(V)$ can be written such as $P(V) := \bigoplus_{n \geq 1} P(V)_n$, where $P(V)_n$ is the k -vector space of products of n elements of V . Moreover, we have $\Delta(x \circ y) := 1 \otimes (x \circ y) + (x \circ y) \otimes 1 + x \otimes (1 \circ y) + y \otimes (x \circ 1)$, for all $x, y \in P(V)$ and $\circ \in P(2)$. The filtration of $P(V)_+$ is then $FrP(V)_+ = k.1 \oplus \bigoplus_{1 \leq n \leq r} P(V)_n$. Therefore, $P(V)_+ := \bigcup_r FrP(V)_+$ and $P(V)_+$ is a connected bialgebra. \square

We will use this theorem to show that there exists a connected Hopf algebra on the augmented free octo-algebra as well as on the augmented free commutative octo-algebra.

Proposition 5.10 *Let $\mathcal{O}(V)$ be the free octo-algebra on a k -vector space V . Extend the binary operations \nwarrow_1 and \searrow_2 to $\mathcal{O}(V)_+$ as follows:*

$$x \nwarrow_1 1 := x, \quad 1 \nwarrow_1 x := 0, \quad 1 \searrow_2 x := x, \quad x \searrow_2 1 := 0, \quad \forall x \in \mathcal{O}(V).$$

In addition, for all operations $\diamond \in \{\searrow_1, \nwarrow_2, \nearrow_1, \nearrow_2, \swarrow_1, \swarrow_2\}$ choose: $x \diamond 1 := 0 = 1 \diamond x$, $\forall x \in \mathcal{O}(V)$. Then,

$$x \prec_1 1 = x, \quad 1 \succ_2 x = x, \quad 1 \vee_1 x := x, \quad x \wedge_2 1 := x, \quad x \bar{\star} 1 = x = 1 \bar{\star} x, \quad \forall x \in \mathcal{O}(V)$$

Moreover, this choice is coherent.

Proof: Observe that we cannot extend \nwarrow_1 and \searrow_2 to the field k , i.e., $1 \nwarrow_1 1$ and $1 \searrow_2 1$ have no sense. Keep notation introduced in that Subsection. Firstly, we claim that this choice is compatible. Let $x, y, z \in \mathcal{O}(V)_+$. We have to show for instance that the relation $(x \nwarrow_1 y) \nwarrow_1 z = x \nwarrow_1 (y \bar{\star} z)$ holds in $\mathcal{O}(V)_+$. Indeed, for $x = 1$ we get $0 = 0$. For $y = 1$ we get $x \nwarrow_1 z = x \nwarrow_1 z$ and for $z = 1$ we get $x \nwarrow_1 y = x \nwarrow_1 y$. We do the same thing with the 27 others and quickly found that the augmented free octo-algebra $\mathcal{O}(V)_+$ is still an octo-algebra.

Secondly, let us show that this choice is coherent. Let $x_1, x_2, x_3, y_1, y_2, y_3 \in \mathcal{O}(V)_+$. We have to show that, for instance:

$$(Eq. [O]_{11}) \quad ((x_1 \otimes y_1) \nwarrow_1 (x_2 \otimes y_2)) \nwarrow_1 (x_3 \otimes y_3) = (x_1 \otimes y_1) \nwarrow_1 ((x_2 \otimes y_2) \bar{\star} (x_3 \otimes y_3)).$$

Indeed, if there exists a unique $y_i = 1$, the other belonging to $\mathcal{O}(V)$, then, by definition we get:

$$(x_1 \bar{\star} x_2 \bar{\star} x_3) \otimes ((y_1 \nwarrow_1 y_2) \nwarrow_1 y_3) = (x_1 \bar{\star} x_2 \bar{\star} x_3) \otimes (y_1 \nwarrow_1 (y_2 \bar{\star} y_3)),$$

which always hold since our choice of the unit action is compatible. Similarly if $y_1 = y_2 = y_3 = 1$. For $y_1 = 1 = y_2$ and $y_3 \in \mathcal{O}(V)$, we get: $0 = 0$, similarly for $y_1 = 1 = y_3$ and $y_2 \in \mathcal{O}(V)$. If $y_1 \in \mathcal{O}(V)$ and $y_2 = 1 = y_3$, the two hand sides of $(Eq. [O]_{11})$ are equal to $(x_1 \bar{\star} x_2 \bar{\star} x_3) \otimes y_1$. As a conclusion, $(Eq. [O]_{11})$ holds in $\mathcal{O}(V) \otimes 1.k \oplus k.1 \otimes \mathcal{O}(V) \oplus \mathcal{O}(V) \otimes \mathcal{O}(V)$. Checking the same thing with the 27 others shows that our choice of the unit action is coherent. \square

Corollary 5.11 *There exists a connected Hopf algebra structure on the augmented free octo-algebra as well as on the augmented free commutative octo-algebra.*

Proof: The first claim comes from the fact that our choice of the unit action is coherent and from Theorem 5.9. For the second remark, observe that our choice is in agreement with the symmetry relations defining a commutative octo-algebra since for instance $x \nwarrow_1^{op} 1 := 1 \searrow_2 x := x$ and $1 \searrow_2^{op} x := x \nwarrow_1 1 := x$, for all $x \in \mathcal{O}(V)$. \square

5.3 Construction of octo-algebras via three pairwise commuting Baxter operators

To produce such exotic associative algebras, it suffices to construct a Baxter operator on a quadri-algebra $(Q, \nearrow, \searrow, \swarrow, \nwarrow)$. By definition, such an operator is a linear map $\xi : Q^{\otimes 2} \rightarrow Q$ such that for all binary operations $\circ \in \{ \nearrow, \searrow, \swarrow, \nwarrow \}$ and all $x, y \in Q$,

$$\xi(x) \circ \xi(y) := \xi(\xi(x) \circ y + x \circ \xi(y)),$$

hold. In particular, if $\star \longrightarrow \nearrow + \searrow + \swarrow + \nwarrow$ denotes the associative operation of Q , then $\xi(x) \star \xi(y) := \xi(\xi(x) \star y + x \star \xi(y))$.

Proposition 5.12 *Let $(Q, \nearrow, \searrow, \swarrow, \nwarrow)$ be a quadri-algebra and $\xi : Q^{\otimes 2} \rightarrow Q$ be a Baxter operator. Define new binary operations as follows: $x \searrow_1 y := x \searrow \xi(y)$, $x \searrow_2 y := \xi(x) \searrow$*

$y, \quad x \nearrow_1 y := x \nearrow \xi(y), \quad x \nearrow_2 y := \xi(x) \nearrow y, \quad x \nwarrow_1 y := x \nwarrow \xi(y), \quad x \nwarrow_2 y := \xi(x) \nwarrow y, \quad x \swarrow_1 y := x \swarrow \xi(y), \quad x \swarrow_2 y := \xi(x) \swarrow y, \text{ for all } x, y \in Q.$

Then, $(Q, \nearrow_1, \nearrow_2, \nwarrow_1, \nwarrow_2, \swarrow_1, \swarrow_2)$ is an octo-algebra.

Proof: Straightforward. Denotes the associative operation of an octo-algebra by $\bar{\star} \longrightarrow \nearrow_1 + \nearrow_2 + \nwarrow_1 + \nwarrow_2 + \swarrow_1 + \swarrow_2$. Fix $x, y, z \in Q$. For instance,

$$\begin{aligned} (x \nwarrow_1 y) \nwarrow_1 z &:= (x \nwarrow \xi(y)) \nwarrow \xi(z), \\ &:= x \nwarrow (\xi(y) \star \xi(z)), \\ &:= x \nwarrow \xi(y \bar{\star} z), \\ &:= x \nwarrow_1 (y \bar{\star} z). \end{aligned}$$

□

Proposition 5.13 *Let β, γ and ξ be three pairwise commuting Baxter operators on an associative algebra (A, μ) . Then, ξ is a Baxter operator on the quadri-algebra $(Q, \nearrow, \nwarrow, \swarrow, \searrow)$ where binary operations are given by:*

$$x \searrow y := \beta\gamma(xy), \quad x \swarrow y := \beta(x)\gamma(y), \quad x \swarrow y := \gamma(x)\beta(y), \quad x \nwarrow y := x\beta\gamma(y).$$

Proof: Fix $x, y \in Q$. For instance, $\xi(x) \searrow \xi(y) := \beta\gamma(\xi(x))\xi(y) = \xi(\beta\gamma(x))\xi(y) = \xi(\beta\gamma(x)\xi(y) + \xi(\beta\gamma(x))y) = \xi(\beta\gamma(x)\xi(y) + \beta\gamma(\xi(x))y) = \xi(x \searrow \xi(y) + \xi(x) \searrow y)$. □

Corollary 5.14 *Let $(A, \mu, \Delta_1 \curvearrowright \Delta_2)$ be a L -circular ϵ -bialgebra, i.e., both (A, μ, Δ_1) and (A, μ, Δ_2) are ϵ -bialgebras and,*

$$(id \otimes \Delta_1)\Delta_2 = (\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1.$$

Then, the three Baxter operators $\beta_1, \gamma_2, \beta_2 : End(A) \rightarrow End(A)$ commute pairwise and the linear map γ_2 is a Baxter operator on the quadri-algebra $End(A)$ whose binary operations are given by:

$$T \searrow S := \beta_1\gamma_2(T)S, \quad T \swarrow S := \beta_1(T)\gamma_2(S), \quad T \swarrow S := \gamma_2(T)\beta_1(S), \quad T \nwarrow S := T\beta_1\gamma_2(S).$$

Proof: Since Δ_2 is coassociative, β_2 and γ_2 commute. Since $(\Delta_1 \otimes id)\Delta_2 = (id \otimes \Delta_2)\Delta_1$, β_1 and γ_2 commute. Since $(id \otimes \Delta_1)\Delta_2 = (id \otimes \Delta_2)\Delta_1$, β_1 and β_2 commute. The corollary holds by applying Proposition 5.13. □

Remark: In addition to the entanglement condition between Δ_1 and Δ_2 , observe the importance of the circular condition for having three pairwise commuting Baxter operators.

Remark: We started with the Baxter operator ξ acting on the quadri-algebra defined by the pair (β, γ) and found eight operations defining an octo-algebra, say O . If the Baxter operator ξ acts on the quadri-algebra defined by the pair (γ, β) , we will find the transpose of the octo-algebra O , see Proposition 5.4. Now, if the Baxter operator γ acts on the quadri-algebra defined

by the pair (β, ξ) , then we will find the octo-algebra of Proposition 5.5, i.e. $Sym(O)_a$. If it acts on the pair (ξ, β) , then we will find the octo-algebra of Proposition 5.6, i.e. $Sym(O)_b$. If the Baxter operator β acts on the quadri-algebra defined by the pair (γ, ξ) , then we will find the octo-algebra of Proposition 5.7, i.e. $Sym(O)_c$. If it acts on the pair (ξ, γ) , we will find the octo-algebra of Proposition 5.8, i.e. $Sym(O)_d$.

Remark: For the time being, except the trivial ones, i.e., Δ_2 proportional to Δ_1 , we do not know any examples of L -circular bialgebras.

5.4 Another construction

Proposition 5.15 *Let (E, \prec_e, \succ_e) , (F, \prec_f, \succ_f) and (G, \prec_g, \succ_g) be three dendriform dialgebras. For all $x, x' \in E$, $y, y' \in F$, $z, z' \in G$, consider the following operations on $E \otimes F \otimes G$:*

$$\begin{aligned}
(x \otimes y \otimes z) \nwarrow_1 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \prec_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \nwarrow_2 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \prec_f y') \otimes (z \succ_g z'), \\
(x \otimes y \otimes z) \searrow_1 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \succ_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \searrow_2 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \succ_f y') \otimes (z \succ_g z'), \\
(x \otimes y \otimes z) \nearrow_1 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \prec_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \nearrow_2 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \prec_f y') \otimes (z \succ_g z'), \\
(x \otimes y \otimes z) \swarrow_1 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \succ_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \swarrow_2 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \succ_f y') \otimes (z \succ_g z').
\end{aligned}$$

Equipped with these eight operations, the k -vector space $E \otimes F \otimes G$ is an octo-algebra. The depth structure is described by the following four operations:

$$\begin{aligned}
(x \otimes y \otimes z) \nwarrow_{12} (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \prec_f y') \otimes (z \star_g z'), \\
(x \otimes y \otimes z) \searrow_{12} (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \succ_f y') \otimes (z \star_g z'), \\
(x \otimes y \otimes z) \nearrow_{12} (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \prec_f y') \otimes (z \star_g z'), \\
(x \otimes y \otimes z) \swarrow_{12} (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \succ_f y') \otimes (z \star_g z').
\end{aligned}$$

The vertical structure is described by the following four operations:

$$\begin{aligned}
(x \otimes y \otimes z) \succ_1 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \star_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \succ_2 (x' \otimes y' \otimes z') &:= (x \succ_e x') \otimes (y \star_f y') \otimes (z \succ_g z'), \\
(x \otimes y \otimes z) \prec_1 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \star_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \prec_2 (x' \otimes y' \otimes z') &:= (x \prec_e x') \otimes (y \star_f y') \otimes (z \succ_g z').
\end{aligned}$$

The horizontal structure is described by the following four operations:

$$\begin{aligned}
(x \otimes y \otimes z) \wedge_1 (x' \otimes y' \otimes z') &:= (x \star_e x') \otimes (y \prec_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \wedge_2 (x' \otimes y' \otimes z') &:= (x \star_e x') \otimes (y \prec_f y') \otimes (z \succ_g z'), \\
(x \otimes y \otimes z) \vee_1 (x' \otimes y' \otimes z') &:= (x \star_e x') \otimes (y \succ_f y') \otimes (z \prec_g z'), \\
(x \otimes y \otimes z) \vee_2 (x' \otimes y' \otimes z') &:= (x \star_e x') \otimes (y \succ_f y') \otimes (z \succ_g z').
\end{aligned}$$


The associative structure is then:

$$(x \otimes y \otimes z) \bar{\star} (x' \otimes y' \otimes z') := (x \star_e x') \otimes (y \star_f y') \otimes (z \star_g z'),$$

as expected.

Proof: Straightforward. □

Example 5.16 Denote by T , the dendriform dialgebra of planar (rooted) binary trees. Consider three copies of T , say T_1 , T_2 and T_3 . Graphically such structures can be represented by placing planar binary trees on a marked directed triangle. On a vertex $i := 1, 2, 3$, a tree from T_i is placed. The operations of the octo-algebra $T_1 \otimes T_2 \otimes T_3$ are then well defined.



Example of planar binary trees on a marked directed triangle.

6 Leibniz algebras and Poisson dialgebras from right Baxter operators

6.1 Theoretical constructions

We end this work on Baxter operators by showing another application to construct particular Leibniz algebras and Poisson dialgebras. Leibniz algebras have been introduced by Loday in [15] to understand periodicity phenomena in K -theory, see also [16]. They are anti-symmetric version of Lie algebras. Similarly, Poisson algebras are the data of a commutative algebra and a Lie bracket verifying compatibility relations. Poisson dialgebras, introduced in [17] are the data of an associative algebra, not necessarily commutative, and a Leibniz bracket verifying compatibility conditions, see also [10]. A way to obtain Leibniz algebras is to start with associative dialgebras, k -vector spaces D equipped with two associative products, \vdash and \dashv , such that for all $x, y, z \in D$,

1. $x \dashv (y \dashv z) = x \dashv (y \vdash z)$,
2. $(x \vdash y) \dashv z = x \vdash (y \dashv z)$,
3. $(x \vdash y) \vdash z = (x \dashv y) \vdash z$.

Associative dialgebras are then Leibniz algebras by defining the bracket $[x, y] := x \dashv y - y \vdash x$, for all $x, y \in D$. This bracket verifies:

$$[[x, y], z] = [[x, z], y] + [x, [y, z]].$$

A way to obtain Poisson dialgebras is to start with associative trialgebras. An associative trialgebra is a k -vector space $(T, \dashv, \vdash, \perp)$ such that (T, \dashv, \vdash) is an associative dialgebra, (T, \perp) an associative algebra and,

1. $(x \dashv y) \dashv z = x \dashv (y \perp z),$
2. $(x \perp y) \dashv z = x \perp (y \dashv z),$
3. $(x \dashv y) \perp z = x \perp (y \vdash z),$
4. $(x \vdash y) \perp z = x \vdash (y \perp z),$
5. $(x \perp y) \vdash z = x \vdash (y \vdash z).$

The operads associated with associative di and trialgebras are Koszul dual respectively to the operads associated with dendriform dialgebras and dendriform trialgebras [17, 4, 9]. We give now a method to construct such types of algebras via right Baxter operators. For that, we define a $\epsilon[R]$ -dialgebra $(A, \mu, \Delta_+, \Delta_-)$ which is a k -vector space such that (A, μ, Δ_+) and (A, μ, Δ_-) are both $\epsilon[R]$ -bialgebras and (A, Δ_+, Δ_-) is a coassociative codialgebra.

Theorem 6.1 *Let $(A, \mu, \Delta_+, \Delta_-)$ be a $\epsilon[R]$ -dialgebra. Then, $(\text{End}(A), \overrightarrow{*}_{\gamma_+}, \overrightarrow{*}_{\gamma_-})$ is an associative dialgebra. Equipped with the dissymmetric commutator:*

$$[T, S] := T \overrightarrow{*}_{\gamma_+} S - S \overrightarrow{*}_{\gamma_-} T,$$

$(\text{End}(A), [\cdot, \cdot])$ is a Leibniz algebra.

Proof: Fix $R, T, S \in \text{End}(A)$ and $a \in A$. The proof can be done without any tedious computations by observing that for computing $\gamma_1(T)\gamma_2(S)(a)$ only the map $(id \otimes \Delta_1)\Delta_2$ is required, but for computing $\gamma_1(T\gamma_2(S))(a)$ only the map $(\Delta_2 \otimes id)\Delta_1$ is needed. For instance, on the one hand,

$$R \overrightarrow{*}_{\gamma_+} (S \overrightarrow{*}_{\gamma_-} T) := R\gamma_+(S\gamma_-(T)).$$

Apply to an element $a \in A$, this equation will only require $(\Delta_- \otimes id)\Delta_+$. On the other hand,

$$(R \overrightarrow{*}_{\gamma_+} S) \overrightarrow{*}_{\gamma_-} T := R\gamma_+(S)\gamma_-(T).$$

Apply to an element $a \in A$, this equation will only require $(id \otimes \Delta_+)\Delta_-$. However, $(\Delta_- \otimes id)\Delta_+ = (\Delta_+ \otimes id)\Delta_-$ since it is an axiom of coassociative codialgebras. Therefore,

$$R \overrightarrow{*}_{\gamma_+} (S \overrightarrow{*}_{\gamma_-} T) = (R \overrightarrow{*}_{\gamma_+} S) \overrightarrow{*}_{\gamma_-} T = R \overrightarrow{*}_{\gamma_+} (S \overrightarrow{*}_{\gamma_-} T).$$

Doing the same thing with the three other equations shows that $(\text{End}(A), \overrightarrow{*}_{\gamma_+}, \overrightarrow{*}_{\gamma_-})$ is an associative dialgebra, with $\overrightarrow{*}_{\gamma_+}$ playing the rôle of \dashv and conversely. \square

Corollary 6.2 *Let $(A, \mu, \Delta_+, \Delta_-, \Delta_\perp)$ be a $\epsilon[R]$ -trialgebra. Then, $(\text{End}(A), \overrightarrow{*}_{\gamma_+}, \overrightarrow{*}_{\gamma_-}, \overrightarrow{*}_{\gamma_\perp})$ is an associative trialgebra. Equipped with the dissymmetric commutator:*

$$[T, S] := T \overrightarrow{*}_{\gamma_+} S - S \overrightarrow{*}_{\gamma_-} T,$$

$(\text{End}(A), [\cdot, \cdot], \overrightarrow{}_{\gamma_\perp})$ is a Poisson dialgebra.*

Proof: The proof is similar to the previous theorem. □

6.2 Constructions from coassociative coverings of directed graphs

Proposition 6.3 *Let V be a non-empty set. Suppose there exist three coproducts Δ_1, Δ_2 and Δ_3 on kV , the free k -vector space spanned by V , verifying co-trialgebras axioms. Extend them on the free associative k -algebra $As(V)$ generated by V as follows,*

$$\Delta_{i,\#}(v_1 \dots v_n) := v_1 \dots v_{n-1} \Delta_i(v_n),$$

for all $i = 1, 2, 3$. Then, $(As(V), \Delta_{1,\#}, \Delta_{2,\#}, \Delta_{3,\#})$ is a $\epsilon[R]$ -trialgebra.

Proof: Straightforward. □

Apply Proposition 6.3 to co-dialgebras and co-trialgebras constructed from the notion of coassociative coverings of directed graphs [10].

Remark: What has been done in this section can be also realised with left versions.

7 Conclusion and opennings

Some mathematical objects can be described with the help of k -vector spaces, say A , equipped with co-operations and associative products verifying compatibility conditions. These give birth to ϵ -bialgebras and their left and right versions. The use of coproducts naturally endow the k -vector space $\text{End}(A)$ with other operations called convolution products. It turns out that studying the placings of operators belonging to $\text{End}(A)$, for instance at the right or the left hand side of such convolution products can be coded through the shifts γ and β and generate special (associative) algebras of a certain type. Therefore the operations coded by these shifts, (or placing operators γ and β and their combinaisons $\alpha\beta, \beta^2$ and so forth) give birth to binary quadratic and non-symmetric operads. For the time being, we do not know neither what the free objects associated with these operads are nor if they are Koszul.

References

- [1] M. AGUIAR. Infinitesimal bialgebras, pre-lie and dendriform algebras. *to appear in "Hopf algebras: Proceedings from an International Conference held at DePaul University"*; *arXiv:math.QA/0211074*.

- [2] M. AGUIAR and J.-L. LODAY. Quadri-algebras. *arXiv:math.QA/0309171*.
- [3] G. BAXTER. An analytic problem whose solution follows from a simple algebraic identity. *Pacific J. Math.*, 10:731–742, 1960.
- [4] K. EBRAHIMI-FARD. Loday-type algebras and the Rota-Baxter relation. *Letters in Mathematical Physics*, 61(2):139–147, 2002.
- [5] B. FRESSE. Koszul duality of operads and homology of partition posets. *Preprint 2002*.
- [6] V. GINZBURG and M. KAPRANOV. Koszul duality for operads. *Duke Math. J.*, 76(1):203–272, 1994.
- [7] S.A. JONI and G.-C. ROTA. Coalgebras and bialgebras in combinatorics. *Stud. Appl. Math.*, 61:93–139, 1979.
- [8] M. KHOVANOV. A categorification of the Jones polynomial. *eprint arXiv:math.QA/9908171*.
- [9] Ph. LEROUX. Ennea-algebras. *eprint, arXiv:math.QA/0309213, (version 2)*.
- [10] Ph. LEROUX. From entangled codipterous coalgebras to coassociative manifolds. *eprint arXiv:math.QA/0301080*.
- [11] Ph. LEROUX. Periodic orbits, coassociative grammar and quantum random walk over \mathbb{Z} . *eprint arXiv:quant-ph/ 0209100*.
- [12] Ph. LEROUX. Tiling the $(n^2, 1)$ -De-Bruijn graph with n coassociative coalgebras. *eprint arXiv:math.QA/ 0209108*.
- [13] Ph. LEROUX. An algebraic framework of weighted directed graphs. *To appear in Int. J. Math. Math. Sci.*, 2003.
- [14] J.-L. LODAY. Scindement d’associativité et algèbres de Hopf.
- [15] J.-L. LODAY. Une version non commutative des algèbres de Lie: Les algèbres de Leibniz. *L’Enseignement Math.*, 39:269–293, 1993.
- [16] J.-L. LODAY. Dialgebras. in *Dialgebras and related operads, Lecture Notes in Math.*, 1763:7–66, 2001.
- [17] J.-L. LODAY and M. RONCO. Trialgebras and families of polytopes. *eprint arXiv:math.QA/0205043*.
- [18] J.-L. LODAY and M. RONCO. Algèbres de Hopf colibres. *C.R.Acad.Sci Paris*, 337(Ser. I):153–158, 2003.
- [19] T. PIRASHVILI. Sets with two associative operations. *Preprint 2003*.
- [20] B. RICHTER. Dialgebren, Doppelgebren und ihre Homologie. *Diplomarbeit, Bonn Universität, unpublished*.
- [21] G.-C. ROTA. Baxter operators, an introduction. in *Gian-Carlo Rota on Combinatorics: Introductory papers and commentaries (J.P.S. Kung, Ed.) Birkhauser, Boston*, pages 504–512, 1995.

- [22] D. VOICULESCU. Free analysis question I: Duality transform for the coalgebra of $\partial_{X:B}$.
arXiv:math.QA/0306172.